

MODEL-BUILDING FOR FRACTIONAL SUPERSTRINGS

KEITH R. DIENES[★]

*Dept. of Physics, McGill University
E. Rutherford Building, 3600 University St.
Montréal, P.Q. H3A-2T8 Canada*

and

S.-H. HENRY TYE

*Newman Laboratory of Nuclear Studies
Cornell University
Ithaca, NY 14853-5001 USA*

ABSTRACT

We investigate heterotic string-type model-building for the recently-proposed fractional superstring theories. We concentrate on the cases with critical spacetime dimensions four and six, and find that a correspondence can be drawn between the new fractional superstring models and a special subset of the traditional heterotic string models. This allows us to generate the partition functions of the new models, and demonstrate that their number is indeed relatively limited. It also appears that these strings have uniquely natural compactifications to lower dimensions. In particular, the $D_c = 6$ fractional superstring has a natural interpretation in four-dimensional spacetime.

[★] E-mail address: dien@hep.physics.mcgill.ca

It is generally accepted that string theory is a realistic hope of unifying all forces and matter, and provides an attractive solution to the problem of quantum gravity and general relativity. To date, the only known consistent theory is the superstring theory^[1] and the closely-related M-theory. One of the significant problems involved in superstring theory is the existence of a multitude of vacua, which these string theories allow. This lack of uniqueness is a problem, since these theories have critical dimension $D = 10$; in order to have a consistent theory in four spacetime dimensions, one must therefore choose a scheme for these six extra dimensions or represent them in terms of arbitrarily chosen additional worldsheet fields. In many ways in which this can be done, resulting in a large number of classical vacua. It is presumed that some dynamical argument might be used to select the vacuum of the physical world, but at present such approaches are not available.

Recently, however, a new approach to superstring model-building has been proposed.^[3] Rather than using the string/heterotic framework, the fundamental principle of string theory called the *fractional superstring* approach is to fix the critical spacetime dimension directly. Such an approach may only serve as a more natural starting-point for

but hopefully also lead to a smaller and thereby more compelling set of self-consistent classical vacua. The basic idea behind the fractional superstring is to replace the traditional right-moving worldsheet supersymmetry found in heterotic strings with a right-moving worldsheet *fractional* supersymmetry parametrized by an integer $K_R \geq 2$. Such a fractional supersymmetry relates worldsheet bosons not to fermions but rather to worldsheet *parafermions*, and one finds that the corresponding critical dimension of such a string theory is

$$D = \begin{cases} 2 + 16/K_R & \text{for } K_R \geq 2 \quad (\text{fractional supersymmetry}), \\ 26 & \text{for } K_R = 1 \quad (\text{no supersymmetry}). \end{cases} \quad (1.1)$$

The case $K_R = 1$ ($D = 26$) corresponds to the traditional bosonic string, and $K_R = 2$ ($D = 10$) corresponds to the traditional super- or heterotic string. The new theories are those for which $K_R > 2$, and we see that by choosing the cases $K_R = 8$ ($D = 4$) or $K_R = 4$ ($D = 6$), we can obtain significantly lower critical dimensions. For these cases we can therefore expect a smaller set of classical vacua and hopefully a more natural description of the physical world.

These fractional superstring theories are indeed natural extensions of the $K_R = 2$ superstring theories, and it is straightforward to relate the two in terms of their underlying worldsheet physics. For the traditional superstring theory, it is well-known that the underlying worldsheet structure is closely related to the $SU(2)_2$ Wess-Zumino-Witten (WZW) theory:^[4] the worldsheet superpartner of the spacetime coordinate X^μ is a Majorana fermion ψ^μ , and this fermionic theory can be simply described by the WZW coset $SU(2)_2/U(1)$. Setting the remaining $U(1)$ boson to an appropriately chosen radius reproduces the full

$SU(2)_2$ WZW theory, yet if we relax the radius, we can interpret this boson as the spacetime coordinate. This decompactification procedure destroys the original worldsheet symmetry of the model, but its superconformal symmetry survives in the coset theory. The *fractional* superstring theory has a similar structure, and is related in precisely the same way to the coset theories for $K \geq 2$. The general coset theories are $SU(2)_K/U(1)$ parafermion theories,^[5,6] and we once again obtain the spacetime field X^μ by decompactifying the remaining WZW boson. The usual supercurrent for $K > 2$ is a new current^[7] (which, equivalently, spin) is $(K+4)/(K+2)$; these new currents transform the traditional spin, and (as we shall see) transform X^μ into a new operator. The energy operator in the parafermion theory is related to this remaining worldsheet symmetry as a new worldsheet symmetry, and to the strings based on these worldsheet symmetries as superstrings. Because the corresponding fractional supersymmetry is non-local on the worldsheet,^[8] the analysis for these theories is typically more involved than for the simpler, local theories. However, as we shall see, concrete progress can indeed be made.

In this paper we shall assume the underlying worldsheet theory of the fractional superstring and concern ourselves primarily with the physical consequences that such fractional string theories allow; other aspects of understanding the underlying Fock-space structure will be discussed in future work.

scattering amplitudes, examining the ghost system and developing a no-ghost theorem, are actively being pursued. Our approach, therefore, is to examine this space of models by studying the allowed partition functions that such models might have; in this way we are able to obtain a number of interesting results.

First, we demonstrate that the fractional superstring partition functions are straightforward generalizations of the traditional $(K_L, K_R) = (2, 2)$ superstring and $(1, 2)$ heterotic string partition functions, and we explicitly develop a general procedure for constructing modular-invariant partition functions for our new models which are consistent with $N = 1$ spacetime supersymmetry.

Second, and perhaps more importantly, by focusing much of our attention on the $(K_L, K_R) = (1, 8)$ and $(1, 4)$ heterotic-type fractional superstring models, we find that we are able to make direct correspondences between these models and the traditional $(1, 2)$ models; these correspondences are possible because both theories are built from identical bosonic left-moving sectors. These correspondences afford us a means of generating what we believe to be valid $(1, K_R)$ -type fractional superstring models, and we present a number of concrete examples with critical spacetime dimensions four and six.

Third, our correspondences suggest that only traditional $(1, 2)$ models with a *maximal* number of spacetime supersymmetries can be related to fractional $(1, K_R)$ models (which were themselves constructed with $N = 1$ spacetime supersymmetry). This result therefore severely constrains the space of fractional superstring models in $D = 4$ and $D = 6$, confirming our expectation that the number of allowed models is indeed relatively small.

Finally, we discuss an intriguing feature with respect to fractional superstring theories: quantum mechanics, local supersymmetry, and together seem to intrinsically select certain “natural” spacetime dimensions in which these fractional superstring theories might be consistent. The “natural” dimensions are even smaller than their traditional counterparts, so that our theories themselves seem to induce supersymmetry in order to achieve Lorentz invariance. These compactifications of traditional string theories, are not at all arbitrary, and the $K = 4$ fractional superstring has a “natural” in four-dimensional spacetime. It turns out that this compactification of spacetime affords us with a means of building models consistent with the fundamental representations of relevant gauge groups.

All of our results therefore not only lend credence to the fractional superstring idea, but may also, we hope, serve as the basis for a systematic model-building program. In particular, the constraints on the models arise from very general principles, and thus should be applicable to all of (K_L, K_R) fractional superstring theories in various spacetime dimensions.

Our goals in this paper are two-fold: not only to provide a more detailed discussion of the ideas discussed above, but we also aim to provide a more complete realization of the original fractional superstring idea than was given in [1]. This paper is somewhat lengthy; its organization is as follows: Section 2 is a self-contained introduction to the fractional

how this approach forms a natural generalization of the traditional superstring approach. We then in Sect. III proceed to survey the algebraic forms we expect partition functions to have for a general (K_L, K_R) string theory: for $K = 1, 2$ we present known models which will play a role in later sections, and for other values of K we introduce the parafermionic string functions^[9] and discuss how they enter into the new total partition functions of (K_L, K_R) models. We also demonstrate that spacetime supersymmetry can be incorporated for these models by choosing these string functions in certain linear combinations, and present a number of important new string-function identities. In Sect. IV we then turn our attention to the heterotic $(1, K_R)$ theories, ultimately deriving various “dictionaries” relating these models to the traditional $(1, 2)$ models. We illustrate the use of these dictionaries by obtaining a number of new fractional-superstring models in $D = 4$ and $D = 6$, and in Sect. V we discuss precisely which traditional models may be “translated” with these dictionaries. In this way we observe an expected truncation in the size of the space of fractional superstring models relative to that corresponding to traditional superstring models in $D < 10$, reflecting the fact that these new models are indeed *in* their critical dimensions. We close in Sect. VI with our discussion of various further issues in fractional superstring model-building, among them the creation of $(1, 4)$ models with chiral fermions and the necessity of compactifying or interpreting the $(1, 4)$ models in four spacetime dimensions. As we will see, these issues are intimately connected, and we expect the dictionaries we derive in Sect. IV to be easily generalizable to these cases as well. In Appendix A we gather together various definitions and properties of the parafermion characters

(or string functions) which play an important role in the construction of the models. In Appendix B we prove an assertion made in Sect. III.

2. FRACTIONAL SUPERSTRINGS

In this section we provide a self-contained introduction to the fractional superstring theory as a natural generalization of the traditional superstring and heterotic string theories. We also review, where necessary, some relevant features of the underlying $\mathbf{Z}_K$ parafermion theories, originally constructed by Zamolodchikov and Fateev.^[5]

As outlined in Sect. I, the basic idea behind the fractional superstring is to modify the worldsheet symmetry in such a manner as to obtain a correspondingly smaller critical spacetime dimension. In order to do this, let us begin by considering the general $SU(2)_K$ WZW theory.^[4] As is well-known, this theory consists of primary fields $\Phi_m^j(z)$ which can be organized into $SU(2)$ representations labelled by an integer j , where $0 \leq j \leq K/2$ and $|m| \leq j$ with $j - m \in \mathbf{Z}$ (for simplicity we are considering only the holomorphic components). Since $SU(2)$ always has a $U(1)$ subgroup which can be bosonized as a free boson φ on a circle of radius $\sqrt{K}$, we can correspondingly factor these primary fields Φ_m^j :

$$\Phi_m^j(z) = \phi_m^j(z) \exp \left\{ \frac{m}{\sqrt{K}} \varphi(z) \right\}. \quad (2.1)$$

Here φ is the free $U(1)$ boson, and the $\phi_m^j(z)$ are the primary fields of the coset $SU(2)_K/U(1)$ theory. This coset theory is the well-known $\mathbf{Z}_K$ parafermion the-

ory,[★] and these fields ϕ_m^j are the corresponding weights (or conformal dimensions):

$$h_m^j = \frac{j(j+1)}{K+2} - \frac{m^2}{K}$$

The fusion rules of these parafermion fields ϕ_m^j theory:

$$[\phi_{m_1}^{j_1}] \otimes [\phi_{m_2}^{j_2}] = \sum_{j=|j_1-j_2|}^r$$

where $r \equiv \min(j_1+j_2, K-j_1-j_2)$ and where the fields ϕ_m^j and their descendants. The character is the Dedekind η -function and the c_n^ℓ are the central charges. These functions will be discussed in more detail. The properties of these functions are collected in Appendix A.

Upon factorizing the primary fields Φ_m^j as in (2.1), the currents factorize as well:

$$\begin{aligned} J^+ &= \sqrt{K} \psi_1 e^{i\varphi} \\ J^0 &= \sqrt{K/2} i \partial \varphi \\ J^- &= \sqrt{K} \psi_1^\dagger e^{-i\varphi} \end{aligned}$$

where the parafermion currents $\psi_i \sim \phi_i^0 \sim \phi_{K-i}^{K/2}$ have conformal dimensions $i(K-i)/K$ in accordance with

★ It is important to realize that for fixed K there can be more than one way to realize the coset theory. Different field contents, where any $[\phi_m^j]$ can appear, are possible. In general, the coupling constants will

see that the parafermion stress-energy tensor $T_{\text{para}}(z) \equiv T_{SU(2)_K}(z) - T_\varphi(z)$ and the parafermion currents ψ_i , $i = 1, 2, \dots, K-1$ form a closed algebra, namely, the $\mathbf{Z}_K$ parafermion current algebra.

Note that ψ_1 acting on a field ϕ_m^j increases the m quantum number by one but does not change the $SU(2)$ spin j . Specifically, we can perform a mode-expansion for ψ_1

$$\psi_1(z) = \sum_{n \in \mathbf{Z}} A_{d+n} z^{-1-d-n+1/K} \quad (2.5)$$

where d is a fractional number and the conformal dimension of A_{d+n} is $-(d+n)$; the $A^\dagger$ mode-expansion for the $\psi_1^\dagger$ field can be handled similarly. In (2.5), of course, the value of d must be chosen appropriately for the particular parafermion field on which $\psi_1(z)$ is to operate, *e.g.*, for consistency we must choose a ψ_1 moding with $d = (2m+1)/K$ when operating on ϕ_m^j . We thus have

$$\begin{aligned} A_{(2m+1)/K+n} : \quad [\phi_m^j] &\rightarrow [\phi_{m+1}^j] \\ A_{(1-2m)/K-n}^\dagger : \quad [\phi_m^j] &\rightarrow [\phi_{m-1}^j] \end{aligned} \quad (2.6)$$

where n are integers. There is also another special field in the parafermion theory, namely the energy operator $\epsilon \equiv \phi_0^1$; operating on a field ϕ_m^j with ϵ preserves the m quantum number but yields sectors with j quantum numbers $j+1$, j , and $j-1$. Specifically, performing a mode-expansion for ϵ as in (2.5) and choosing d as indicated below, we find the actions of its modes:

$$\begin{aligned} \epsilon_{-2(j+1)/(K+2)+n} : \quad [\phi_m^j] &\rightarrow [\phi_m^{j+1}] \\ \epsilon_n : \quad [\phi_m^j] &\rightarrow [\phi_m^j] \\ \epsilon_{2j/(K+2)+n} : \quad [\phi_m^j] &\rightarrow [\phi_m^{j-1}] . \end{aligned} \quad (2.7)$$

Of course, (2.6) and (2.7) are valid only when p is even. We thus see that the two fields $\psi_1 = \phi_1^0$ and ϵ generate the entire set of fields $[\phi_m^j]$, starting from any one field.

The next step in the fractional superstring construction is to decompactify the free boson φ , replacing it with a circle of infinite radius; this field X will be interpreted as the time coordinate. This decompactification produces a theory underlying $SU(2)$ WZW symmetry of our theory that remains.

In fact, the symmetry remaining after the bosonization is more than simple conformal symmetry. We can see this in the following way. Let us construct the current $\hat{J}$ [7]

$$\hat{J} \equiv \epsilon \partial X + : \epsilon \varphi :$$

where $\epsilon(z)$ is the energy-operator field and where $: \epsilon \varphi :$ is a normal-ordered product (and is in fact a parafermion descendent of ψ_1). For $K \geq 2$, of course, the field ϵ has conformal dimension

$$\Delta_\epsilon = \frac{2}{K+2}$$

and it can be shown that the normal-ordered term $: \epsilon \varphi :$ has conformal dimension $\Delta_{:\epsilon\varphi:} = 1 + \Delta_\epsilon$. This is a non-trivial fact, implying

$:\epsilon\epsilon:$ appears in the $\epsilon\epsilon$ operator product expansion as follows:

$$\epsilon(z)\epsilon(w) = \frac{1}{(z-w)^{2\Delta_\epsilon}} + \dots + (z-w)^{1-\Delta_\epsilon} : \epsilon\epsilon : (w) + \dots \quad (2.10)$$

Thus, for $K \geq 2$ the current $\hat{J}$ in (2.8) has conformal dimension (or equivalently, spin):

$$\Delta_{\hat{J}} = 1 + \frac{2}{K+2} = \frac{K+4}{K+2}. \quad (2.11)$$

On the worldsheet this current $\hat{J}$ forms a closed algebra^[8] with $T(z)$, where $T(z) \equiv T_X(z) + T_{\text{para}}(z)$ is the stress-energy tensor of the decompactified boson field X plus that of the $\mathbf{Z}_K$ parafermion theory. Thus we see that $\hat{J}$ indeed generates an additional worldsheet symmetry which we refer to as a *fractional* worldsheet supersymmetry. Note that for $K > 2$ the dimensions Δ_ϵ and $\Delta_{\hat{J}}$ are not simple half-integers. Thus, our underlying worldsheet $(\hat{J}, T)$ algebra is non-local, with Riemann *cuts* (rather than poles) appearing in the various OPEs. Note that this $(\hat{J}, T)$ algebra is merely the simplest algebra that can be constructed. For other fractional-superstring applications, this $(\hat{J}, T)$ algebra can indeed be extended to include additional currents.

In order to achieve a sensible interpretation for a D -dimensional space-time, we associate the decompactified bosonic field X with a single spacetime coordinate and tensor together D copies of the (X, ϕ_m^j) (or boson plus $\mathbf{Z}_K$ parafermion) theory. We therefore obtain the fractional supersymmetry current

$$J = \epsilon^\mu \partial X_\mu + : \epsilon^\mu \epsilon_\mu : \quad (2.12)$$

where the Lorentz indices $\mu = 0, 1, \dots, D-1$ are to be contracted with the

Minkowski metric. A crucial issue, however, is the choice of dimension D for arbitrary $K > 2$. A formal derivation of D is not yet established path: since the central charge of the

$$c_0 = 1 + \frac{2K-2}{K+2} =$$

one would need only to determine the central charge of the *fractionally-super*reparametrization ghosts, and then require that this quantity cancels this quantity. Such a calculation is not yet established. The present remains to be done. Another approach is to determine the dimension of this theory and find a dimension D and a value of K for which extra null states appear; such work is not yet done. One can take a third approach^[3] and require that the model produced have a sensible phenomenology, e.g. a massless graviton in the case of a closed string theory and a massless vector particle in the case of an open string theory. This is a *requirement* and not an assumption, for we require only those fractional string theories which conform to this requirement. are, from this standpoint, phenomenologically viable. This requirement is not difficult to implement.

For a general closed string theory, the graviton and dilaton both arise from the same right-moving excitations. The difference is in that they are tensored with dissimilar left-movers. Thus our requirement simply becomes a requirement on the left-movers.

demand that this state exist (*i.e.*, satisfy the physical-state conditions) and be massless. For a graviton or spacetime vector particle in a string theory with arbitrary $K \geq 2$, this state is

$$|V\rangle_R = \zeta_\mu \epsilon_{-2/(K+2)}^\mu |p\rangle_R \quad (2.14)$$

where $|p\rangle_R$ is the right-moving vacuum state with momentum p , ζ_μ is a polarization vector, and where $\epsilon_{-2/(K+2)}^\mu$ is the lowest excitation mode of the parafermion field ϵ . Note that the moding of the ϵ field follows from its conformal dimension. This state is indeed the analogue of what appears for the $K = 2$ case, in which a single lowest-mode excitation of a worldsheet Neveu-Schwarz fermion produces the needed state: $|V\rangle_R = \zeta_\mu \epsilon_{-1/2}^\mu |p\rangle$ where $|p\rangle$ is the usual Neveu-Schwarz vacuum state and $\epsilon_{-1/2}^\mu = b_{-1/2}^\mu$ is the lowest Neveu-Schwarz creation operator. (For the bosonic string the ϵ field is absent, and we accordingly substitute the bosonic creation operator a_{-1}^μ for $b_{-1/2}^\mu$ or $\epsilon_{-2/(K+2)}^\mu$. The same argument then applies.) Thus, requiring the state (2.14) to be massless, we find in general that the vacuum state $|p\rangle$ in (2.14) must have vacuum energy $VE = -v = -2/(K+2)$. This information can also be stated in terms of the fractional superstring character $\chi(q)$. In general the character has a q -expansion of the form

$$\chi(q) = \sum_n a_n q^n \quad (2.15)$$

where $q \equiv \exp(2\pi i\tau)$, τ is the complex modular parameter of the torus, and where a_n is the number of propagating degrees of freedom at mass level $M^2 = n$.

We thus see that the fractional superstring character

$$\chi(q) = q^{-v} (1 + \dots) = q^{-2/(K+2)}$$

where inside the parentheses all q -powers are non-negative integers, is

The only remaining non-trivial physical-state condition

is

$$J_{2/(K+2)} |V\rangle_R = 0$$

where J_n are the modes of the fractional supercurrent. This condition can indeed be shown^[3] to yield the expected total number of physical states on the polarization vector ζ_μ (which is consistent with the polarization vector of the massless vector particle). The state therefore has only $D - 2$ polarizations (or degrees of freedom). It follows that at all mass levels of the physical spectrum, the number of dimensions worth of polarization states are precisely $D - 2$. Note that this latter assertion is *not* an ad hoc assumption; it follows directly from modular invariance. We can also see that the longitudinal and time-like components of the physical states (the removal of the corresponding q^0 terms in the partition function) (as we will see in Sect. III) comprise the total number of physical functions. However, the string functions c_n^ℓ , under the action of T modular transformations, form an admissible representation of the modular group. Modular invariance therefore requires

string functions from the total fractional-superstring partition functions, which in turn implies that there is indeed a large string gauge symmetry whose gauge-fixing provides the physical conditions effectively striking out all states involving longitudinal and/or time-like modes from every mass level in the theory.

However, we now recall that conformal invariance requires the character $\chi(q)$ to take the form

$$\chi(q) = q^{-c/24} (1 + \dots) \quad (2.18)$$

where c is the total effective conformal anomaly of the propagating degrees of freedom. Since each spacetime dimension contributes the central charge c_0 given in (2.13), and since we have determined that effectively only $D - 2$ dimensions contribute to propagating fields, we have $c = (D - 2)c_0$. Thus, comparing (2.16) and (2.18) and substituting (2.13), we find the result

$$D = 2 + \frac{16}{K}, \quad K \geq 2. \quad (2.19)$$

We see, then, that the critical dimension of the fractional superstring is a function of the level K of the $SU(2)_K/U(1)$ coset (*i.e.*, of the $\mathbf{Z}_K$ parafermionic theory): for $K = 2$ we have $D = 10$, for $K = 4$ we have $D = 6$, and for $K = 8$ we obtain $D = 4$. For $K = 1$ we need only set $v = 1$ in (2.16) [as explained after (2.14)], whereupon the argument above yields $D = 26$. Thus, by appropriately choosing K and building the corresponding worldsheet fractional supersymmetry as discussed above, we can hope to obtain a series of new string theories with a variety of critical dimensions.

It is easy to check that the $K = 1$ and $K = 2$ theories are the bosonic string and superstring respectively. For $K = 1$, the theory is bosonized with a single boson φ ; therefore, the “parafermion” theory) is trivial, containing only the identity. There are therefore no partners for our bosonic fields X^μ (which they become), and hence for $K = 1$, the theory is conformal symmetry. This reproduces, of course, the critical dimension of the bosonic string. Similarly, for $K = 2$, the theory (the $\mathbf{Z}_2$ parafermion theory) is simply the Ising model, with the identity ϕ_0^0 (the identity), $\phi_0^1 = \phi_1^0$ (which is of course the parafermion current ψ_1 , or equivalently the Ising spin field σ), and $\phi_{-1/2}^{1/2}$ (the conjugate spin field). The worldsheet spin fields are responsible for the critical dimension; together these fields form a Ramond worldsheet algebra. For $K = 2$, the identity and ψ fields: when appropriately mixed, form a worldsheet field whose excitations ultimately produce the critical dimension. We see that the field content of the $SU(2)_2/U(1)$ coset worldsheet Majorana fermion theory, and the bosonic X^μ fields is an ordinary worldsheet theory. This reproduces the traditional superstring, and the tensoring of a $K = 1$ and $K = 2$ theory reproduces the critical dimension. Thus, that this fractional superstring language provides a way to classify and uniformly handle all of the traditional string theories, doing it also points the way to their non-trivial extensions.

3. PARTITION FUNCTIONS FOR FRACTIONAL SUPERSTRINGS

Having thus presented the underlying basis for the fractional superstring theories, we turn our attention to the partition functions that these theories must have. We begin by reviewing the generic forms of these partition functions, and start with the known $K = 1$ and $K = 2$ cases to establish our notation and conventions. We will see, once again, that the $K > 2$ string theories have partition functions whose forms are straightforward extensions of those of the traditional cases. This will therefore permit us to write the partition functions for all of the (K_L, K_R) theories we shall consider in a common language.

3.1. TRADITIONAL STRING THEORIES

The first case to consider, of course, is the pure bosonic string; since there is no left-moving or right-moving worldsheet supersymmetry, we may refer to this in our new fractional superstring language as the $(K_L, K_R) = (1, 1)$ case. As discussed in Sect. II, the critical spacetime dimension for this string theory is $D = 26$, and thus this theory contains 26 bosonic worldsheet fields X^μ , each of which contributes to the total one-loop partition function a factor

$$\text{each boson} \implies \frac{1}{\sqrt{\tau_2} \eta \bar{\eta}} . \quad (3.1)$$

Here $\eta(\tau)$ is the well-known Dedekind η -function, and $\tau \equiv \tau_1 + i\tau_2$ is the torus modular parameter. Note that this factor is explicitly real because the left-

and right-moving components of each boson contribute equally. (We have already seen that only $D - 2 = 24$ transverse degrees of freedom are physical (as is evident in a light-cone gauge approach), our $(1, 1)$ string is therefore of the form

$$\mathcal{Z}_{(1,1)} = \tau_2^{-12} \frac{1}{|\eta|^{48}} = \tau_2^{-12} \Delta^{-24}$$

where we have defined $\Delta \equiv \eta^{24}$. Note that the weight k , which we will denote k , is in general given by

$$k = 1 - D/2$$

where D is the number of spacetime dimensions. For the bosonic string, $D = 26$, so $k = -12$. Hence, for the $(1, 1)$ string in $D = 26$ dimensions, we have agreement with (3.2). In the theory of modular forms, η is known as the modular *weight*; the η -function is a modular function of weight k and $T : \tau \rightarrow \tau + 1$ as a modular function of weight k . The holomorphic and anti-holomorphic pieces of the partition function have weight $k = -12$. Modular-invariance for the entire partition function requires the modular weight of its holomorphic factors equals the weight of its anti-holomorphic factors, and that this weight also equal the pole weight. We can see that in (3.2) this is indeed the case.

The next string to consider is the traditional superstring. It has a full superconformal worldsheet symmetry. In the left-moving sectors, in our new language this is simply the $(1, 1)$ string.

That its critical dimension is 10 follows from (2.19), or equivalently from the traditional argument in which the conformal anomaly $c = -26 + 11 = -15$ from the reparametrization ghosts and their superpartners must be cancelled by the contributions from D worldsheet boson/fermion pairs, each of which has total central charge $c_0 = 1 + \frac{1}{2} = \frac{3}{2}$ [in accordance with (2.13) for $K = 2$]. In light-cone gauge only the eight transverse field pairs propagate, and therefore this theory has a total worldsheet field content consisting of eight real bosons (each with a left-moving and right-moving component) and eight Majorana fermions (or eight Majorana-Weyl left-movers and eight Majorana-Weyl right-movers). These fermions, we recall, can be described in our new viewpoint as simply the fields of the $SU(2)_2/U(1)$ Ising model, and we are free to group pairs of these Majorana-Weyl fermions together to form single complex Weyl fermions. The factor contributed by each of the eight bosons to the one-loop partition function is given in (3.1), and the contribution of each complex Weyl fermion is

$$\text{each Weyl fermion} \implies \frac{\vartheta}{\eta} \quad (3.4)$$

where ϑ represents one of the well-known Jacobi ϑ -functions defined in Appendix A.* Since the contributions of right-moving fields are the complex-conjugates of those of left-moving fields, the total partition function for the

* Which particular ϑ -function is appropriate depends on the boundary conditions assigned to the fermion as it traverses the two cycles of the torus. Only periodic or antiperiodic boundary conditions will yield the Jacobi ϑ -functions, but if a fermion is chosen to have periodic/periodic boundary conditions then its zero-modes cause the total partition function to vanish identically.

(2,2) string therefore takes the form

$$\mathcal{Z}_{(2,2)} \sim \tau_2^{-4} \frac{1}{|\eta|^{16}} \sum \left| \frac{\vartheta}{\eta} \right|^8 \sim \tau_2^{-4} \Delta$$

where ϑ^4 indicates four (not necessarily identical) ϑ -functions. The summation in (3.5) is over fermion boundary conditions in order to achieve modular invariance. In this section we use the schematic notation in (3.5), employing the general forms of partition functions. Note that Δ transforms under the modular group with weight $k = -4$, and the η -factors in (3.5) — a holomorphic and anti-holomorphic factors in (3.5) — are such that $k = -4$ for this theory. This result is only consistent with spacetime dimension $D = 10$ according to (3.3).

We shall be most concerned in this paper with the traditional heterotic string has a right-moving sector as in the superstring; hence the right-moving fields consist, as before, of eight bosons and eight fermions, and thus in order to cancel the $c = 26$ conformal anomaly from the reparametrization ghosts we must additionally have eight left-moving fermions (or equivalently 16 complex Weyl fermions, or 32 real Weyl fermions ϕ). Our traditional heterotic string is therefore

partition function of the form

$$D = 10: \quad \mathcal{Z}_{(1,2)} \sim \tau_2^{-4} \Delta^{-1} \overline{\Delta}^{-1/2} \sum \vartheta^{16} \overline{\vartheta}^4. \quad (3.6)$$

Note that we continue to have $k = -4$ from each factor in this partition function, in agreement with (3.3). As examples of (3.6) which will be relevant later, we can look at those known $(1,2)$ $D = 10$ theories (or models) which have spacetime supersymmetry. As is well-known,^[2] there are only two such self-consistent models: these have gauge groups $SO(32)$ and $E_8 \otimes E_8$. Their partition functions respectively are as follows:

$$\begin{aligned} D = 10, SO(32): \quad \mathcal{Z} &= \left(\frac{1}{2}\right)^2 \tau_2^{-4} \mathcal{K} (\beta^4 + \gamma^4 + \delta^4) \\ D = 10, E_8 \otimes E_8: \quad \mathcal{Z} &= \left(\frac{1}{2}\right)^3 \tau_2^{-4} \mathcal{K} (\beta^2 + \gamma^2 + \delta^2)^2 \end{aligned} \quad (3.7)$$

where we have made the following definitions which we will use throughout:

$$\begin{aligned} \beta &\equiv \vartheta_2^4, \quad \gamma \equiv \vartheta_3^4, \quad \delta \equiv \vartheta_4^4, \\ J &\equiv \gamma - \beta - \delta, \\ \mathcal{K} &\equiv \Delta^{-1} \overline{\Delta}^{-1/2} \overline{J}. \end{aligned} \quad (3.8)$$

Note that the partition functions (3.7) are indeed of the form (3.6), with the holomorphic parts factorizing into group characters to reflect the underlying group structure. The quantity J in (3.8) is of course the Jacobi factor, and the spacetime supersymmetry of these models (or equivalently, the vanishing of their partition functions) arises from the identity $J = 0$. It is in fact a further

identity that

$$\beta^4 + \gamma^4 + \delta^4 = \frac{1}{2} (\beta^2 + \gamma^2 + \delta^2)^2$$

as a consequence of which the two partition functions are equal without use of the Jacobi identity $J = 0$.

It is straightforward to check the modularity of the partition functions (3.7). For any $A \equiv \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in the modular group, the so-called “stroke” operator $[A]: f \rightarrow f[A]$ where

$$(f[A])(\tau) \equiv (c\tau + d)^{-k} f\left(\frac{a\tau + b}{c\tau + d}\right)$$

(here k is the modular weight of f). It then follows from the transformation properties of the η and ϑ functions (see

$$\begin{aligned} \beta[S] &= -\delta, \quad \beta[T] = \gamma, \\ \gamma[S] &= -\gamma, \quad \gamma[T] = \delta, \\ \delta[S] &= -\beta, \quad \delta[T] = \beta, \\ \mathcal{K}[S] &= +\mathcal{K}, \quad \mathcal{K}[T] = -\mathcal{K} \end{aligned}$$

Thus, $\mathcal{K}$ is itself modular-invariant (*i.e.*, invariant under S and T). In each partition function the other factors involved are themselves modular-invariant as well.

It is also possible to construct self-consistent models for $D < 10$; one needs simply, for example, to re-introduce extra degrees of freedom or in some other manner higher

(*e.g.*, through compactification). We stress, however, that all such methods do not alter the underlying critical dimension away from 10; these theories remain theories of the $(1,2)$ variety. For example, one method of constructing $(1,2)$ models in $D = 4$ involves fermionization: each of the six extra left- or right-moving bosonic degrees of freedom can, for example, be represented in terms of two free worldsheet Majorana-Weyl fermions (or a single free Weyl fermion).^[10] The propagating field content for this four-dimensional $(1,2)$ theory therefore consists of two left-moving and two right-moving bosons, as well as 22 left-moving and 10 right-moving Weyl fermions. The total partition function for such a $D = 4$ $(1,2)$ theory hence takes the form

$$D = 4: \quad \mathcal{Z}_{(1,2)} \sim \tau_2^{-1} \Delta^{-1} \bar{\Delta}^{-1/2} \sum \vartheta^{22} \bar{\vartheta}^{10}. \quad (3.12)$$

Note that once again (3.3) is satisfied, with $k = -1$ for $D = 4$. As an example, we present the partition function of what is possibly the simplest such spacetime-supersymmetric model that can be constructed. Named Model *M1* in Ref. [10], it has gauge group $SO(44)$, the largest possible in $D = 4$; its (modular-invariant) partition function is

$$D = 4, SO(44): \quad \mathcal{Z} = \left(\frac{1}{2}\right)^2 \tau_2^{-1} \mathcal{K} \left(|\beta| \bar{\beta} \beta^5 + |\gamma| \bar{\gamma} \gamma^5 + |\delta| \bar{\delta} \delta^5 \right). \quad (3.13)$$

Since the “maximal” gauge group allowed in D dimensions is $SO(52 - 2D)$, we see that this model is the $D = 4$ analogue of the $D = 10$ $SO(32)$ model in (3.7).

We note for future reference that this idea can in fact be generalized to obtain $(1,2)$ -type theories in any spacetime dimension $D \leq 10$. We in general

obtain the partition function form

$$\text{general } D: \quad \mathcal{Z}_{(1,2)} \sim \tau_2^k \Delta^{-1}$$

where k is given by (3.3) and

$$n = 26 - D = 24$$

$$\bar{n} = 14 - D = 12$$

Much of our work will be concerned with models with $k = -2$, $n = 20$, and $\bar{n} = 8$. The “maximally symmetric” $D = 6$ supersymmetric model with $SO(40)$ gauge group has gauge group $SO(40)$; its partition function is $\mathcal{Z} = (\frac{1}{2})^2 \tau_2^{-2} \mathcal{K} \times (\bar{\beta} \beta^5 + \bar{\gamma} \gamma^5 + \bar{\delta} \delta^5)$. For the $D = 6$ supersymmetric models we will be discussing,

$$D = 6, SO(40): \quad \mathcal{Z} = \left(\frac{1}{2}\right)^2 \tau_2^{-2} \mathcal{K}$$

$$D = 6, SO(24) \otimes E_8: \quad \mathcal{Z} = \left(\frac{1}{2}\right)^3 \tau_2^{-2} \mathcal{K} \times (\bar{\beta} \beta^5 + \bar{\gamma} \gamma^5 + \bar{\delta} \delta^5)$$

$$D = 6, SO(24) \otimes SO(16): \quad \mathcal{Z} = \left(\frac{1}{2}\right)^3 \tau_2^{-2} \mathcal{K} + \bar{\gamma} \gamma^2 (\beta^3 + \gamma^3 + \delta^3)$$

We again observe the familiar factorization, with $\mathcal{Z} = \mathcal{Z}_{(1,2)} \mathcal{Z}_{(2,2)}$ in (3.7). It is simple, using (3.11), to see that the partition function is modular-invariant.

3.2. FRACTIONAL SUPERSTRING THEORIES

The partition functions for *fractional* superstrings can be determined in precisely the same manner as above. Since we know the critical dimension D for a given value of K , we can readily deduce the forms that partition functions must take for general (K_L, K_R) combinations. The factor contributed to the total partition function from each worldsheet boson is given, as before, by (3.1); recall that this factor includes the contributions from both holomorphic and anti-holomorphic (or left- and right-moving) components. The factor contributed by each worldsheet parafermion, however, is a generalization of (3.4); in general we have

$$\text{each parafermion} \implies \eta c \quad (3.17)$$

(the above is for left-moving parafermions; right-moving parafermions contribute the complex-conjugate). Here η is the usual Dedekind function, and c schematically represents one of the so-called parafermionic *string functions*.^[6,9] These functions are defined in Appendix A, but for our present purposes we need record only the following facts. These functions c_n^ℓ may be defined in terms of q -expansions which depend on K as well as the two parameters ℓ and n , and one string function c_{2m}^{2j} may correspond to more than one parafermion field ϕ_m^j (for example, the distinct fields ϕ_m^j and ϕ_{-m}^j both give rise to c_{2m}^{2j}). Each c_n^ℓ is an eigenfunction under $T : \tau \rightarrow \tau + 1$, and under $S : \tau \rightarrow -1/\tau$ they mix forming a closed set; furthermore, they transform under the S and T transformations with the *negative* modular weight $k = -\frac{1}{2}$.

As expected, these string functions (which are the generalizations of the traditional modular functions for the $K=1$ case) are indeed precisely what occurs. For the $K=1$ case, the string function is the function c_0^0 , and since our $SU(2)_1/U(1)$ theory has only one parafermion ϕ_0^0 , we quickly have

$$K=1 : \quad \eta c_0^0 = 1 \implies$$

Thus, for $K=1$ the one string function c_0^0 is the Dedekind function. Similarly, for $K=2$ there are precisely three string functions, and these are related to the three fermionic Jacobi theta functions.

$$K=2 : \quad \begin{cases} 2(c_1^1)^2 \\ (c_0^0 + c_0^2)^2 \\ (c_0^0 - c_0^2)^2 \end{cases}$$

For $K > 2$, of course, the string functions involve more than one parafermion.

Therefore, looking first at the (K_L, K_R) case, (these are the generalizations of the usual Type I and Type II theories) the worldsheet field content consists (in light-cone gauge) of $16/K$ transverse coordinate bosons and $16/K$ each of left- and right-moving parafermions. We therefore obtain the form for the total (K, K) partition function

$$\mathcal{Z}_{(K,K)} \sim \tau_2^{-8/K} \sum \bar{c}^1$$

Note that $k \equiv -8/K = 1 - D/2$ in accordance with the general formula.

the η -functions have cancelled between the bosonic and parafermionic contributions. Thus, in our cases of interest we obtain:

$$\begin{aligned} D=6: \quad \mathcal{Z}_{(4,4)} &\sim \tau_2^{-2} \sum \bar{c}^4 c^4 \\ D=4: \quad \mathcal{Z}_{(8,8)} &\sim \tau_2^{-1} \sum \bar{c}^2 c^2 \end{aligned} \quad (3.21)$$

Similarly, one can consider the analogues of heterotic string theories; these would be the $(K_L, K_R) = (1, K)$ situations. The right-moving sectors of these theories are precisely as in the (K, K) cases, and hence the anti-holomorphic parts of their partition functions have the same form as in (3.20). Since their left-moving sectors must already contain the left-moving components of the coordinate bosons, we must augment their left-moving field contents to achieve conformal anomaly cancellation. The contribution of the bosons to the central charge is of course $2 + 16/K$, and since this sector is a bosonic theory we must cancel the usual ghost contribution $c = -26$. This requires additional matter fields with central charge

$$\Delta c = 8 \left(\frac{3K-2}{K} \right), \quad (3.22)$$

and we may choose these fields to be Δc complex Weyl worldsheet fermions [each of which has the partition-function contribution given in (3.4)]. Thus, for a general heterotic-type $(1, K)$ theory we expect a partition function of the form

$$\begin{aligned} \mathcal{Z}_{(1,K)} &\sim \tau_2^{-8/K} \sum \bar{c}^{16/K} \left(\frac{1}{\eta} \right)^{16/K} \left(\frac{\vartheta}{\eta} \right)^{8(3K-2)/K} \\ &\sim \tau_2^{-8/K} \sum \bar{c}^{16/K} \Delta^{-1} \vartheta^{8(3K-2)/K}. \end{aligned} \quad (3.23)$$

Note that since $\Delta c = 8(3K-2)/K = n$ [where n is the quantity defined in

(3.15)], the holomorphic part of this partition function is as we obtained in (3.14) for general spacetime dimension D to be expected; these left-moving sectors are bosonic. The difference, however, is that the heterotic $(1, K)$ theories exist for *all* values of K .

For completeness, we should also note that in the case, to consider $(1, K)$ theories in spacetime dimension D , for example, choose to retain D of the above bosons and $D_c - D$ bosons with as many worldsheet Weyl fermions as such a sector the partition function contribution

$$\left(\frac{1}{\eta} \right)^{D-2} (\eta c)^{16/K} \left(\frac{\vartheta}{\eta} \right)^{D_c-D}$$

the first factor is the contribution of the $D-2$ bosons that of the original $D_c-2 = 16/K$ parafermions and the fermionized $D_c - D$ bosons. Thus, for example, in the $D=4$ theory: one can consider a $(1, 8)$ theory in which two dimensions are in some way compactified. We shall discuss this latter possibility in Sect. 4.

In fact, use of these parafermionic string theories together in a simple way *all* of the partition functions for the general (K_L, K_R) theory in arbitrary spacetime dimension. For a given value of K_R , we have considered the two cases of traditional string theories as well as our new

simply

$$\mathcal{Z}_{(K_L, K_R)} = \tau_2^k \sum_i L_i^{(K_L)} \overline{R_i^{(K_R)}} \quad (3.25)$$

where the $L_i^{(K_L)}$ are the contributions from the left-moving sectors and the $R_i^{(K_R)}$ are from the right-movers. In the above formula k is always given by (3.3) (where D is the dimension of spacetime, not the critical dimension). If we define the critical dimensions $D_c^R = 2 + 16/K_R$ and $D_c^L = 2 + 16/K_L$ (or 26 if the corresponding $K = 1$), then the general forms for the L_i and R_i can be given as follows:

$$L_i^{(K_L)} \sim c^{D_c^L - 2} \vartheta^{D_c^L - D}, \quad R_i^{(K_R)} \sim c^{D_c^R - 2} \vartheta^{D_c^R - D}. \quad (3.26)$$

We can easily check the special cases $K = 1$ and $K = 2$. For a (1,1) theory formulated in arbitrary $D \leq D_c = 26$, (3.18) yields $L^{(1)} = R^{(1)} = \Delta^{-1} \vartheta^{26-D}$; for $D = D_c$ this therefore reproduces (3.2), and for $D < D_c$ this reproduces the holomorphic part of (3.14) for the (1,2) theory. Similarly, for $K_R = 2$ in arbitrary dimension $D \leq D_c$, (3.19) yields $R^{(2)} = \Delta^{-1} \vartheta^{14-D}$, in accord with the anti-holomorphic part of (3.14). Thus, (3.25) and (3.26) are indeed the most general partition function forms for the traditional as well as the fractional string theories, brought together in a natural way through our use of the parafermionic string functions.

In order to construct sensible partition functions having the above forms, it is first necessary to find suitable combinations of the string functions which can replace the c^{D_c-2} factors above. There are several requirements. First, we want

linear combinations which are tachyon-free: the

$$\sum c^{D_c-2} = \sum_n c_n$$

we must have $a_n = 0$ for all $n < 0$. Second, the spectrum of massless particles: this means the corresponding to a massless sector in the linear combinations $a_0 \neq 0$. Third, since we would like to use the linear combinations in heterotic partition functions, we must demand modular-invariance under $T^4 : \tau \rightarrow \tau + 4$ (this is because the expressions with which we will be dealing should be modular-invariant). In the language of (3.27), this means that $a_n = 0$ for all $n \neq 0$. Finally, for a given K , we require that the linear combinations each involve $D_c - 2$ powers of c . For $K \geq 2$ this means that we require $16/K$ string functions, and we require 24.

It is clear that linear combinations satisfying these requirements exist for the $K = 1$ and $K = 2$ cases; for $K = 1$

$$A_1 = (c_0^0)^{24} = \Delta$$

★ This expression A_1 actually has $a_n \neq 0$ for $n < 0$, a known result that the bosonic string contains tachyons.

and for $K = 2$ we can choose:

$$\begin{aligned} A_2 &= 8(c_0^0)^7 c_0^2 - 8(c_1^1)^8 + 56(c_0^0)^5 (c_0^2)^3 + 56(c_0^0)^3 (c_0^2)^5 + 8c_0^0 (c_0^2)^7 \\ &= \frac{1}{2} \Delta^{-1/2} J . \end{aligned} \quad (3.29)$$

We note that in each of these cases only one linear combination is necessary to achieve closure under S and T , and we observe that $A_2 = 0$ as a consequence of the Jacobi identity $J = 0$. This is simply a reflection of the *spacetime* supersymmetry of the superstring. These two expressions have, of course, already been determined: $\mathcal{Z}_{(1,1)}$ in (3.2) is merely $\tau_2^{-12} |A_1|^2$, and $\tau_2^{-4} |A_2|^2$ is merely a special (spacetime-supersymmetric) case of $\mathcal{Z}_{(2,2)}$ in (3.5).

Similarly, it is possible to construct linear combinations satisfying all of the above constraints for the $K = 4$ and $K = 8$ cases as well.^[3] These are as follows. For $K = 4$ we have the two combinations:

$$\begin{aligned} A_4 &= 4(c_0^0 + c_4^4)^3 (c_0^2) - 4(c_2^2)^4 + 32(c_4^4)^3 (c_2^2) - 4(c_0^2)^4 , \\ B_4 &= -4(c_2^2)^2 (c_0^2)^2 + 8(c_0^0 + c_4^4) (c_0^2) (c_4^4)^2 \\ &\quad + 4(c_0^0 + c_4^4)^2 (c_2^2) (c_4^4) \end{aligned} \quad (3.30)$$

and for $K = 8$ we have the three combinations:

$$\begin{aligned} A_8 &= 2(c_0^0 + c_8^8) (c_0^2 + c_0^6) - 2(c_4^4)^2 + 8(c_4^8 c_4^6) - 2(c_0^4)^2 , \\ B_8 &= 4(c_0^0 + c_8^8) (c_4^6) + 4(c_0^2 + c_0^6) (c_4^8) - 4(c_0^4 c_4^4) , \\ C_8 &= 4(c_2^2 + c_2^6) (c_2^8 + c_6^8) - 4(c_2^4)^2 . \end{aligned} \quad (3.31)$$

[In fact, combinations satisfying these requirements exist for the $K = 16$ ($D = 3$) case as well.^[11]] The transformation properties of these functions under

S and T will be given in Sect. IV; in particular, A_2 has a q -expansion of the form $q^h(1 + \dots)$, where the exponents are non-negative integers and where $h = 1/2$ for B_4 and B_8 , and $h = 3/4$ for C_8 . Thus, only particles can contribute to these partition functions. For A_8 : indeed, the terms $(c_0^0)^{D_c-3} c_0^2$ can be interpreted as spacetime vector particles, and the terms $-(c_0^0)^{D_c-3} c_0^6$ as corresponding to massless spacetime fermions. We will, therefore, take these combinations as the basis for constructing our fractional superstring partition functions for the c^{D_c-2} factors in the $K = 4$ and $K = 8$ cases.

In fact, it turns out that there is one additional constraint on the expressions in (3.30) and (3.31). Just as the ϑ -function identity $J = \gamma - \beta - \delta = 0$, it can be shown that the parafermionic string-function expressions vanish when

$$A_4 = B_4 = A_8 = B_8 = C_8 = 0 .$$

Thus, for $K = 4$ we have the two new Jacobi identities, and for $K = 8$ we have the three new identities. As we shall see, these exist for the $K = 16$ case as well.^[11] This result can then serve as the mechanism by which any fractional superstring can be made consistent with spacetime supersymmetry. Spacetime supersymmetry must therefore have

these Jacobi-like factors, and the $K = 2$ full Jacobi identity $A_2 = 0$ appears merely as the $K = 2$ special case of the more general identities for general even integers K .

We emphasize that such B and C sectors, necessitated by closure under $[S]$ for the $K = 4$ and $K = 8$ theories, are completely new features arising only for the $K > 2$ fractional superstrings, and as such they are responsible for much of the new physics these strings contain. For example, not only do they lead (as we have seen) to *multiple* independent Jacobi identities, but they may also be responsible for *self-induced* compactifications of these strings. We will discuss this possibility in Sect. VI.

In fact, the Jacobi identity is not the only famous identity which generalizes to higher K . For $K = 2$, there is another well-known identity involving the η and ϑ -functions:

$$\vartheta_2 \vartheta_3 \vartheta_4 = 2\eta^3 ; \quad (3.33)$$

this identity relates the fermion characters ϑ_i to the boson character η , and hence we may refer to this as a bosonization identity. It turns out^[11] that (3.33) is only the $K = 2$ special case of another series of identities, each relating the more general $\mathbf{Z}_K$ parafermion characters c_n^ℓ for a different $K \geq 2$ to the boson character η . In fact, there also exist various *other* series of identities whose $K = 2$ special cases are known and have well-understood physical interpretations; proofs of all of these identities, as well as methods for generating them for arbitrary K , are presented in Ref. [11]. We therefore see that the

string functions c_n^ℓ provide a uniquely compelling way to generalize previously known string-theory results, prove new ones, and other identities whose physical interpretation has yet to be explored. Thus we see again (this time on the level of the formalism) by leading directly to these string-function expressions that the fractional superstring construction does indeed provide a natural generalization of the traditional string theories.

4. DICTIONARIES FOR MODEL-BUILDING

Having established that spacetime supersymmetry can be incorporated by using the expressions (3.30) and (3.31), we now turn our attention to the construction of actual supersymmetric (K_L, K_R) models. We focus our attention primarily on the heterotic $(K_L, K_R) = (1, 4)$ and $(1, 8)$ cases, and construct a procedure for generating models in these classes. Our procedure involves “translating” or drawing correspondences between the $(1, K)$ models and known $(1, 2)$ models in $D = 2 + 16/K$, and we construct “dictionaries” which enable these translations to take place for a given K . We find that these dictionaries are intuitive and practical, and furthermore (as we will see in Sect. V) they yield substantially and understandably smaller spaces of $(1, K)$ models than $(1, 2)$ models in $D < 10$. In particular, we will find that only those $(1, 2)$ models which have a maximal number of spacetime supersymmetries are translatable; these are the models with $N = N_{\max}$ SUSY, where

$$N_{\max} = \begin{cases} 1 & \text{for } D = 10 \\ 2 & \text{for } D = 6 \\ 4 & \text{for } D = 4 \end{cases} \quad (4.1)$$

We should point out that throughout this and the next section we will be focusing our attention on those $(1, 2)$ models whose partition functions can be built from Jacobi ϑ -functions, in accordance with the presentation in Sect. III. This is indeed a broad class of models, but it is not all-inclusive. However, the dictionaries we will be developing for these models rest on very general principles, and we expect this dictionary idea to be equally applicable to other methods of $(1, 2)$ model-construction as well.

4.1. MODULAR INVARIANCE

As discussed in Sect. III, models of the $(1, K)$ type in dimensions $D = 2 + 16/K$ must have partition functions

$$\mathcal{Z}_{(1,K)} \sim \tau_2^k \sum \bar{c}^{16/K} \Delta^{-1}$$

and spacetime supersymmetry can be incorporated by taking the values $\overline{A_K}$, $\overline{B_K}$, and (for $K = 8$) $\overline{C_K}$ slightly, and refer to these quantities collectively as $\overline{F}$. For $K = 8$ and $i = 1, 2$ for $K = 4$. Supersymmetric partition functions

$$\mathcal{Z}_{(1,K)} = \tau_2^k \sum_i \overline{A}_{(K)}^i$$

where the F 's take the forms

$$F_{(4)}^i \sim \Delta^{-1} \sum \vartheta^{20} \quad \text{and} \quad F_{(8)}^i \sim \Delta^{-1} \sum \vartheta^{20}$$

Let us now determine the constraints we impose on the partition functions by modular-invariance. It follows from the transformation properties of the individual string functions c_n^ℓ under the $[S]$ -transform that the expressions $A_{(K)}^i$ transform as

$$A_{(K)}^i[S] = e^{4\pi i/K} \sum_j U_{ji} A_{(K)}^j$$

where the matrices $M_{(K)}$ are

$$M_{(4)} = \begin{pmatrix} 1/2 & 3 \\ 1/4 & -1/2 \end{pmatrix}, \quad M_{(8)} = \begin{pmatrix} 1/2 & 1/2 & 1 \\ 1/2 & 1/2 & -1 \\ 1/2 & -1/2 & 0 \end{pmatrix}, \quad (4.6)$$

and where the stroke operator $[S]$ was defined in (3.10). Demanding $[S]$ -invariance of the total partition function $\mathcal{Z}$ therefore yields

$$\mathcal{Z}[S] = e^{-4\pi i/K} \tau_2^k \sum_{i,j} M^{ij} \overline{A^j} F^i[S] \equiv \tau_2^k \sum_j \overline{A^j} F^j, \quad (4.7)$$

or

$$F^i = e^{-4\pi i/K} \sum_j (M^t)^{ij} F^j[S] \quad (4.8)$$

where M^t is the transpose of M . Since $M^2 = (M^t)^2 = \mathbf{1}$ for both cases $K = 4$ and 8, we can immediately solve (4.8) for $F^j[S]$, yielding the constraint

$$F^i[S] = e^{4\pi i/K} \sum_j (M^t)^{ij} F^j. \quad (4.9)$$

Similarly, under $[T]$ the A 's have the transformation

$$A_{(K)}^i[T] = \sum_j N_{(K)}^{ij} A_{(K)}^j \quad (4.10)$$

where the matrices $N_{(K)}$ are

$$N_{(4)} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad N_{(8)} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -i \end{pmatrix}. \quad (4.11)$$

Proceeding precisely as above, we find the add

$$F^i[T] = \sum_j N^{ij}$$

which in this case imply simply that each F^i must transform as does the corresponding A^i . This is also clear under $T : \tau \rightarrow \tau + 1$ means that in $\mathcal{Z} = \tau_2^k \sum_m$ only if $m - n \in \mathbf{Z}$. Since each A^i has a q -expansion (where inside the parentheses all powers of q are non-negative) the corresponding F^i must take the same form with the content of (4.12).

We now give a general procedure for obtaining operators P_ℓ which satisfy both (4.9) and (4.12). In the $K = 4$ case, after which the $K = 8$ case will be straightforward, we consider the space $\mathcal{F}$ of polynomials in t (so that $f[T^4] = f$ for all $f \in \mathcal{F}$), and establish that the operators P_ℓ ($\ell = 0, \dots, 3$) in this space. These operators P_ℓ

$$P_\ell f \equiv \frac{1}{4} \sum_{n=0}^3 \exp\left\{-\frac{i\pi n}{4}\right\} f\left(\frac{t - \omega^n}{1 - \omega^n}\right)$$

where $f \in \mathcal{F}$ is any polynomial in this space. We have $\sum_\ell P_\ell = \mathbf{1}$ and $P_\ell P_{\ell'} = P_\ell \delta_{\ell\ell'}$. These operators select out the ℓ -th Fourier mode when operating on any $f \in \mathcal{F}$, P_ℓ selects out the

f which have powers equal to $\ell/4$ modulo unity – *i.e.*,

$$(P_\ell f)[T] = \exp\left\{\frac{i\pi\ell}{2}\right\} (P_\ell f) \quad \forall f \in \mathcal{F} . \quad (4.14)$$

For example, if $f = \vartheta_2^2 \vartheta_3^4 \vartheta_4^8$, then

$$\begin{aligned} P_1 f &= \tfrac{1}{2} \vartheta_2^2 (\vartheta_3 \vartheta_4)^4 (\vartheta_3^4 + \vartheta_4^4) , \\ P_3 f &= \tfrac{1}{2} \vartheta_2^2 (\vartheta_3 \vartheta_4)^4 (\vartheta_4^4 - \vartheta_3^4) , \\ P_0 f &= P_2 f = 0 . \end{aligned} \quad (4.15)$$

Let us also define (for the sake of typographical convenience) the operator $\mathcal{S} \equiv \exp\{-4\pi i/K\}[S]$. Then our procedure is as follows. Choose an $f \in \mathcal{F}$ of the form ϑ^{26-D} , and calculate the quantity

$$\mathbf{f} \equiv \Delta^{-1} \lim_{n \rightarrow \infty} \left[\tfrac{1}{2} (1 - P_1)(1 + \mathcal{S}) \right]^n f . \quad (4.16)$$

If $P_3 \mathcal{S} P_3 \mathbf{f} = 0$, then up to one common scale factor the corresponding solution for the F 's is:

$$F^1 = P_0 \mathcal{S} P_3 \mathbf{f} , \quad F^2 = -P_2 \mathcal{S} P_3 \mathbf{f} , \quad F^3 = P_3 \mathbf{f} . \quad (4.17)$$

It is easy to see how this procedure works. The goal is to construct a set of F 's which are distinguished by their eigenvalues under $[T]$, and which furthermore are closed under $[S]$. This we achieve by constructing a quantity $\mathbf{f}$ which itself is $\mathcal{S}$ -invariant, and from which the individual F 's can be obtained by projection. The crucial element in $K = 8$, however, is the fact that this $\mathbf{f}$

must satisfy $P_1 \mathbf{f} = 0$; we cannot accommodate equalling $1/4$ modulo unity in building the desired solution, since there is no corresponding term D_8 with this property. The difficulty as follows. Starting from any f , a first guess for the quantity $\mathbf{f}^{(0)} = \tfrac{1}{2} \Delta^{-1} (1 + \mathcal{S}) f$. We then enforce $P_1 \mathbf{f}^{(0)} = 0$ by modifying the guess: $\mathbf{f}^{(1)} = (1 - P_1) \mathbf{f}^{(0)}$. However, $\mathbf{f}^{(1)}$ is $\mathcal{S}$ -invariant, and we therefore re-apply the process iterates until we have finally achieved a quantity $\mathbf{f}$ with no P_1 -projection – *i.e.*, which is preserved under $\tfrac{1}{2}(1 + \mathcal{S})$ and $(1 - P_1)$. We write this solution for $\mathbf{f}$ as $\mathbf{f} = \mathbf{f}^{(0)} + \mathbf{f}^{(1)}$ of course if this iterative process fails to converge. (In practice, however, with f restricted to the form ϑ^{26-D} , it converges almost immediately.) Having thus found $\mathbf{f}$, we then define F^1, F^2, F^3 as above. Since the $i = 3$ component of (4.9) tells us that

$$\mathcal{S} F^3 = M_{(8)}^{13} F^1 + M_{(8)}^{23} F^2$$

we must correspondingly define $F^1 = P_0 \mathcal{S} F^3$ and $F^2 = -P_2 \mathcal{S} F^3$. These choices are consistent only if $P_3 \mathcal{S} F^3 = 0$, which is true since $M_{(8)}^{33} = 0$ in (4.6).

Generating solutions for the $K = 4$ case is more complicated. The space $\mathcal{F}$ of polynomials to be that generated by ϑ^{26-D} for all $f \in \mathcal{F}$, and consequently we need consider only the operators P_0 and P_2 (since $P_1 f = P_3 f = 0$ for all $f \in \mathcal{F}$).

$f \in \mathcal{F}$, we define

$$\mathbf{f} \equiv \frac{1}{2} \Delta^{-1} (1 + \mathcal{S}) f , \quad (4.19)$$

whereupon we quickly have the solutions (up to a common scale factor)

$$F^1 = P_0 \mathbf{f} , \quad F^2 = 4 P_2 \mathcal{S} P_0 \mathbf{f} . \quad (4.20)$$

Note that there were two reasons this $K = 4$ case was significantly simpler than the $K = 8$ case. First, for $K = 4$ we must demand $P_1 \mathbf{f} = P_3 \mathbf{f} = 0$ instead of the more difficult $K = 8$ constraints $P_1 \mathbf{f} = 0$, $P_3 \mathbf{f} \neq 0$; the presence of *two* zero-constraints in $K = 4$ instead of only one allowed us to subsume them together into a restriction in the space $\mathcal{F}$. Second, the matrix $M_{(4)}$ has no zero entries; hence the F 's can always be found by simple projections and only their relative normalizations need be determined. In $K = 8$, however, we must further assert $P_3 \mathcal{S} P_3 \mathbf{f} = 0$; as stated above, this occurs because $M_{(8)}^{33} = 0$.

Given these procedures for generating solutions for the F 's satisfying (4.9) and (4.12) in the $K = 4$ and $K = 8$ cases, it is easy to build modular-invariant partition functions of the proper forms. Let us first construct some examples for the $(1, 4)$ case. Taking $f = 2\beta^5$ [where we remind the reader of the definitions in (3.8)], we find $\mathbf{f} = \Delta^{-1}(\beta^5 + \delta^5)$, whereupon (4.20) yields the results

$$F^1 = \frac{1}{2} \Delta^{-1} (\gamma^5 + \delta^5) , \quad F^2 = \Delta^{-1} [2\beta^5 + (\gamma^5 - \delta^5)] . \quad (4.21)$$

Since the internal gauge symmetry for such models is determined, as usual, by the left-movers, we can quickly identify this solution as corresponding to gauge

group $SO(40)$, the largest allowed in six space dimensions. For example, we start with $f = 2\beta^3(\beta^2 + \gamma^2 + \delta^2)$: the solution is $\mathbf{f} = \Delta^{-1}(\beta^3 + \gamma^3 + \delta^3)$ and the solutions

$$F^1 = \frac{1}{2} \Delta^{-1} (\gamma^3 + \delta^3) (\beta^2 + \gamma^2 + \delta^2) , \quad F^2 = \Delta^{-1} (\beta^3 + \gamma^3 + \delta^3) (\gamma^2 - \delta^2) + 2\beta^2 (\gamma^3 - \delta^3) ,$$

The presence of the E_8 character $\beta^2 + \gamma^2 + \delta^2$ corresponds to this solution as $SO(24) \otimes E_8$. For $K = 8$, we can choose $f = \gamma^2(\beta^3 + \gamma^3 + \delta^3)$. From this

$$\begin{aligned} F^1 &= \Delta^{-1} \left\{ \frac{1}{2} (\gamma^3 + \delta^3) (\gamma^2 + \delta^2) + \beta^2 (\gamma^3 - \delta^3) \right\} , \\ F^2 &= \Delta^{-1} \left\{ (\gamma^3 + \delta^3) (\gamma^2 - \delta^2) + \beta^2 (\gamma^3 - \delta^3) \right\} , \end{aligned}$$

which (as we will see) can be identified with the E_8 character.

One can similarly construct partition functions for other values of K . For instance, starting with $f = 2\beta^{11/2}$, we find (or rather, we find $\mathbf{f} = \Delta^{-1}(\beta^{11/2} + \delta^{11/2})$, which immediately leads to

$$\begin{aligned} F^1 &= \frac{1}{2} \Delta^{-1} (\gamma^{11/2} + \delta^{11/2}) , \\ F^2 &= \frac{1}{2} \Delta^{-1} (\gamma^{11/2} - \delta^{11/2}) , \\ F^3 &= \Delta^{-1} \beta^{11/2} . \end{aligned}$$

Note that $P_3 \mathcal{S} P_3 \mathbf{f} = 0$, so this solution is invariant under the E_8 character. The solution corresponds to $SO(44)$, the largest allowed in six space dimensions.

4.2. DICTIONARIES FOR MODEL-CONSTRUCTION

While thus far it has been quite straightforward to identify the gauge groups corresponding to our partition functions, we have been dealing only with the simplest of cases; furthermore, in principle almost any properly-chosen function f can serve in generating solutions, and we require a way to discern which of all possible solutions correspond to bona-fide fractional superstring *models*. Toward this end we now develop a method for generating partition functions which we believe do precisely this in the heterotic $(1, K)$ cases, and for which the underlying physics is substantially more transparent. Our approach rests on two fundamental observations.

The first observation has to do with the existence of a model with maximal gauge symmetry for the $(1, 2)$ heterotic string in D dimensions. For any value $D \leq 10$, there is always a self-consistent $(1, 2)$ model which can be formulated with gauge group $SO(52 - 2D)$ – such a model is, in a sense, the starting point in model-building, for all other models can be obtained from it by altering this known solution via orbifolding, *e.g.*, by adding twists, altering the boundary conditions of worldsheet fields, adding new sectors, *etc.*, all of which tends to break the gauge group and correspondingly add new terms to the partition function. We therefore assert that such a self-consistent model exists as well for the $(1, K)$ fractional superstring, and has gauge group $SO(52 - 2D) = SO(48 - 16/K)$; indeed, the validity of this assertion follows directly from the (assumed) self-consistency of the $K > 2$ fractional superstring right-moving sector and the near-decoupling of the gauge sector (the $K = 1$ left-

moving sector), as will be discussed below. Since the unique partition functions corresponding to these two models are given by (4.23) and (4.24), it therefore follows that these two models correspond to actual $(1, K)$ models.

Our second and more important observation has to do with the relationship between the left- and right-moving sectors of a model. It is well-known for the traditional $(1, 2)$ heterotic string that the left-moving sector carries with it all of the information concerning spacetime symmetries and particle representations; the right-moving sector carries with it the linkage to spacetime physics, including spacetime supersymmetry. One builds a model by choosing the left- and right-fermionic sectors so that certain physical constraints are maintained: worldsheet (super)conformal invariance, modular invariance, incorporates proper level-matching, spacetime Lorentz invariance, and physically sensible internal (GSO-like) projections (ensuring consistent spacetime spin-statistics). Of course, not all of these requirements are independent. Some of these requirements can be checked by examining the left- and right-moving sectors separately: among these are worldsheet (super)conformal invariance, spacetime Lorentz invariance, and modular invariance. Modular invariance, on the other hand, must be checked jointly. Thus, when building a model, one must satisfy all of these constraints: those which involve the left- and right-moving sectors independently (guaranteeing that they are each

those (*i.e.*, modular invariance and the implied level-matching) which insure that they are properly coupled or linked to each other.

In the case of the heterotic $(1, K)$ fractional superstring, we expect the same situation to prevail: we must determine those $K = 1$ left-moving sectors which are themselves internally self-consistent, and then we must join them with our (assumed self-consistent) $K > 2$ right-moving sector in such a way that modular-invariance is satisfied. Fortunately, the underlying physics of a $K = 1$ left-moving sector is well-understood; for example, descriptions of it in terms of lattices, orbifolds, or Fock-space spectrum-generating formulae abound. In particular, it is well-known how to construct valid $(1, 2)$ models which satisfy *all* of the physical model-building constraints we have listed. Therefore, one might hope to be able to build valid $(1, K)$ models by first building valid $(1, 2)$ models, and then “replacing” their $K = 2$ right-moving sectors with our new $K > 2$ right-moving sectors in such a way that modular-invariance (the sole “linking” constraint) is not violated. Such a procedure would thereby guarantee, in the language of the previous subsection, a set of F^i ’s which themselves are known to correspond to valid $K = 1$ left-moving sectors.

It turns out that these arguments can be phrased directly in terms of a correspondence or “dictionary” between right-moving $K = 2$ physics and $K > 2$ physics, in the sense that they may be substituted for each other in this way when building models. At the level of the partition function (which has been the basis of our approach), this means that we are able to draw a correspondence between the respective Θ -functions of these right-moving sectors. For the $K >$

2 theory these Θ -functions are simply the expressions we have been using to build our partition functions, and for $K = 2$ these functions are the usual Jacobi ϑ -functions. We will now construct a dictionary relating the expressions for $K > 2$ and $K = 2$ in the sense that the two underlying sectors can be used interchangeably in building self-consistent models of $(1, K)$ type.

It is fairly straightforward to construct the Θ -functions and the ϑ -functions, for our first assumption [the solutions (4.21) and (4.24)] allows us to make the assumption of maximal gauge symmetry. Let us first concern ourselves with the fact from Sect. III that the supersymmetric $(1, 2)$ model with gauge group $SO(40)$ has the partition function

$$\mathcal{Z}_{SO(40)} = \left(\frac{1}{2}\right)^2 \tau_2^{-2} \mathcal{K}(\overline{\beta}\beta^5)$$

(where $\mathcal{K} \equiv \Delta^{-1} \overline{\Delta}^{-1/2} \overline{\mathcal{J}}$). Repeatedly making

$$Aa + Bb = \frac{1}{2}(A + B)(a + b) +$$

it turns out that we can write

$$\begin{aligned} \overline{\beta}\beta^5 + \overline{\gamma}\gamma^5 + \overline{\delta}\delta^5 &= \frac{1}{4}(\overline{\beta} + \overline{\gamma} - \overline{\delta})(\beta + \gamma + \delta) \\ &+ \frac{1}{4}(\overline{\beta} - \overline{\gamma} + \overline{\delta})(\beta - \gamma + \delta) \\ &+ \frac{1}{2}(\overline{\gamma} + \overline{\delta})(\gamma - \delta) \end{aligned}$$

We therefore have

$$\begin{aligned} \mathcal{Z}_{SO(40)} = & \frac{1}{4} \tau_2^{-2} \Delta^{-1} \overline{\Delta}^{-1/2} \overline{J} \left\{ \frac{1}{4} (\overline{\beta} + \overline{\gamma} - \overline{\delta}) [2\beta^5 + (\gamma^5 - \delta^5)] \right. \\ & \left. + \frac{1}{2} (\overline{\gamma} + \overline{\delta}) (\gamma^5 + \delta^5) - \frac{1}{4} \overline{J} [2\beta^5 - (\gamma^5 - \delta^5)] \right\} \end{aligned} \quad (4.28)$$

for the $(1, 2)$ maximally symmetric model in $D = 6$.

The next step involves properly interpreting some of the factors in (4.28). Since this is the partition function of a $(1, 2)$ model, the critical dimension is 10, and reducing the spacetime dimension to 6 through fermionization (as discussed in Sect. III) yields worldsheet matter consisting of the 4 transverse bosonic coordinate fields, 16 Majorana-Weyl right-moving fermions, and 40 Majorana-Weyl left-moving fermions. This is why each term in (4.28) contains two anti-holomorphic powers of $\overline{\beta}$, $\overline{\gamma}$, or $\overline{\delta}$ (recall that $J \equiv \gamma - \beta - \delta$), and five holomorphic powers of β , γ , or δ . Of these 16 right-moving fermions, four are the superpartners of the bosonic coordinate fields (and hence carry spacetime Lorentz indices) while the remaining twelve are internal and carry only internal quantum numbers. However, of these twelve, four had previously been the superpartners of the (now fermionized) $10 - 6$ coordinate bosons, and as such their degrees of freedom (in particular, their toroidal boundary conditions) must be chosen to be the same as those of the four fermions carrying Lorentz indices. In Ref. [10], for example, this is the result of the so-called “triplet” constraint, which arises by demanding the periodicity or anti-periodicity of the worldsheet supercurrent on the worldsheet torus. Thus, the 16 right-moving Majorana-Weyl fermions split into two groups: the first eight (which include the four

with spacetime Lorentz indices) must all have the same boundary conditions, and the remaining eight (all of which are internal) must have different boundary conditions independently. As we can see from the partition function, the eight right-moving worldsheet fermions are responsible for the factor of $\overline{J}$ to produce the overall factor of $\overline{J}$: this is indeed the result of spacetime supersymmetry. The remaining eight right-moving fermions are internal, and these are responsible for the remaining terms in (4.28), the $\overline{\delta}$ within the braces in (4.28) above.

This understanding is very important, for it explains the origin of the factor in the second line of (4.28). As we stated earlier, the factor of $\overline{J}$ in the first line of (4.28) is a consequence of spacetime supersymmetry: the identity $\overline{J} = 0$ represents the cancellation of spacetime bosonic states against spacetime fermionic states. The factor of $\overline{J}$, however, arises from exclusively internal fermions, and thus for this term the identity $\overline{J} = 0$ represents the cancellation between particles of the *same* spacetime (and hence the same particle states). Thus, the last term in (4.28) is not necessary whatsoever, and may be legitimately dropped.

$$\begin{aligned} \mathcal{Z}_{SO(40)} = & \frac{1}{4} \tau_2^{-2} \Delta^{-1} \overline{\Delta}^{-1/2} \overline{J} \times \\ & \times \left\{ \frac{1}{4} (\overline{\beta} + \overline{\gamma} - \overline{\delta}) [2\beta^5 + (\gamma^5 - \delta^5)] \right\} \end{aligned}$$

for this $(1, 2)$ model, or equivalently

$$\mathcal{Z}_{SO(40)} = \tau_2^{-2} \left\{ \overline{R_1^{K=2, D=6}} L_1^{SO(40)} + \dots \right\}$$

where the two factors arising from the right-moving $K = 2$, $D = 6$ theory are

$$\begin{aligned} R_1^{K=2,D=6} &\equiv \frac{1}{4} \Delta^{-1/2} J(\gamma + \delta) \\ R_2^{K=2,D=6} &\equiv \frac{1}{16} \Delta^{-1/2} J(\beta + \gamma - \delta) , \end{aligned} \quad (4.31)$$

and where the two factors arising from the left-moving $K = 1$ $SO(40)$ theory are:

$$\begin{aligned} L_1^{SO(40)} &\equiv \frac{1}{2} \Delta^{-1} (\gamma^5 + \delta^5) \\ L_2^{SO(40)} &\equiv \Delta^{-1} (2\beta^5 + \gamma^5 - \delta^5) . \end{aligned} \quad (4.32)$$

Now that we have rewritten (4.25) in the form (4.30), we can compare this result with the partition function for the $(1, 4)$ $SO(40)$ model. Recall that this latter partition function was found to be [see (4.3) and (4.21)]:

$$\begin{aligned} \mathcal{Z} &= \tau_2^{-2} \left\{ \frac{1}{2} \overline{A}_4 \Delta^{-1} (\gamma^5 + \delta^5) + \overline{B}_4 \Delta^{-1} [2\beta^5 + (\gamma^5 - \delta^5)] \right\} \\ &= \tau_2^{-2} \left\{ \overline{A}_4 L_1^{SO(40)} + \overline{B}_4 L_2^{SO(40)} \right\} ; \end{aligned} \quad (4.33)$$

note that one can verify the overall normalization of (4.33) by counting the numbers of low-lying (*e.g.*, tachyonic or massless fermionic) degrees of freedom. It is clear that (4.30) and (4.33) have the same left-moving holomorphic pieces $L_i^{SO(40)}$, and therefore we can relate them to each other, *i.e.*,

$$\mathcal{Z}_{(1,2)} \stackrel{*}{=} \mathcal{Z}_{(1,4)} \quad \text{for } SO(40) \quad (4.34)$$

(where $\stackrel{*}{=}$ indicates this relation or correspondence), if we make the following

correspondences:

$$\begin{aligned} A_{(4)}^1 &\equiv A_4 \stackrel{*}{=} R_1^{K=2,D=6} \\ A_{(4)}^2 &\equiv B_4 \stackrel{*}{=} R_2^{K=2,D=6} \end{aligned}$$

where the R 's are given in (4.31). This result is the “dictionary” relating A_4 and B_4 to the Jacobi theta functions (which will interchange their corresponding $K = 2$ and $D = 6$ labels at the partition-function level) in order to build the $(1, 4)$ model. We remark that in principle it is just as straightforward to build the $(2, 4)$ model suitable for building $(2, 4)$ models; indeed, the partition function we have obtained can themselves be taken as the partition function from the self-consistent *left-moving* sectors of the $(2, 4)$ model.

Note that if we had *not* dropped the theta function (4.28), we would have required a *third* theta function to relate to

$$R_3^{K=2,D=6} \equiv \Delta^{-1/2} J(\gamma - \beta - \delta)$$

Here we have explicitly indicated the origins of the various factors, labelling with subscripts whether they arise from spacetime or internal degrees of freedom. The string-function expression C_4 does not exist. The GSO-projection $J_{\text{int}} = 0$ enables us to avoid this in an internally self-consistent manner, and thereby

There are several things to note about this dictionary. First, we observe that this dictionary is self-consistent as a relation between modular functions, with each side of the relation transforming under the modular group with weight $k = -2$ [as appropriate for $D = 6$, according to (3.3)]. Furthermore, the expressions on the right side of (4.35) mix under the $[S]$ -transformation with the same matrix $M_{(4)}$ found for the $A_{(4)}^i$, and the same, of course, holds true for the $[T]$ -transformation and the $N_{(4)}$ matrix.[★] Indeed, the right sides of (4.35) each have q -expansions of the forms $q^h(1+\dots)$ where inside the parentheses all q -powers are integral and where h -values are equal (mod 1) on both sides of each equation. Additionally, this dictionary incorporates the spacetime supersymmetry of A_4 and B_4 in a natural way, allowing these expressions (which are themselves the “Jacobi identities” for $K = 4$) to correspond to expressions proportional to the spacetime factor J , the $K = 2$ Jacobi identity.

In fact, from this dictionary it is now clear that one cannot expect to factorize the string-function expressions $A_{(4)}^i$ into two pieces, one of which might vanish on its own. In the $(1,2)$ theory [*i.e.*, the right side of (4.35)], the J factor is the result of spacetime-related degrees of freedom, and the remaining factors linear in β , γ , and δ in (4.35) are the contributions from the internal right-moving degrees of freedom. For the $K = 4$ theory, however, we are *in* the critical dimension, and hence *all* right-moving modes carry spacetime information and play a part in yielding spacetime supersymmetry. We therefore do not

★ One must not forget that when performing the $[S]$ transformation on the right sides of (4.35), self-consistency requires setting factors of J_{int} to zero.

expect to be able to factorize the left sides of (4.35). The dictionary also provides correspondences between individual factors on the right side of (4.35).

This dictionary allows us to easily generate new models: we start with a known $(1,2)$ model corresponding to a certain spacetime dimensions, and then make the “translation” to the $K = 4$ right-moving sector. As examples, we consider two-dimensional $(1,2)$ models whose partition functions are proportional to $SO(24) \otimes E_8$ model is particularly simple to translate. In this case the analogue of (4.27) becomes

$$\bar{\beta}\beta^3 + \bar{\gamma}\gamma^3 + \bar{\delta}\delta^3 = \frac{1}{4}(\bar{\beta} + \bar{\gamma} - \bar{\delta})[2\beta^3 + (\gamma^3 - \delta^3)]$$

where we have set $\bar{J}_{\text{internal}} = 0$. Upon substituting this into the internal partition function in (3.16) and translating back to the original variables, we find the partition function (4.3) with the F ’s given by β, γ, δ . This confirms that the solution in (4.21) corresponds to the $SO(24) \otimes E_8$ (left-moving) gauge group claimed. Similarly translating the partition function in (3.16), we obtain the solution for the $SO(24) \otimes E_8$ partition function, confirming the interpretation of that solution as corresponding to the $SO(24) \otimes E_8$ with the quoted gauge group.

We stress again, of course, that this dictionary does not duplicate the results found earlier, for the general case. The previous subsection merely assures the creation of solutions $\mathcal{Z}$. It is the crucial fact that we can derive the dictionary from the

our dictionary translation which guarantees their interpretation as the partition functions of actual models. To illustrate this fact, let us consider the creation of $(1, 4)$ models having gauge groups G of the form $G_{12} \otimes G_8$, where G_r denotes a simple (and simply-laced) group of rank r . Recall that the procedure given in the previous subsection allows us to specify our possible $(1, 4)$ functions $\mathcal{Z}$ in terms of the simpler $\mathcal{S}$ -invariant expression $\mathbf{f}$. Since we expect a gauge group G_r to reveal itself in $\mathcal{Z}$ through the presence of factors which are linear combinations of β^p , γ^p , and δ^p where $p = r/4$ [for example, the factor (or character) corresponding to E_8 is $(\beta^2 + \gamma^2 + \delta^2)$], we can survey many such $\mathcal{Z}$'s by building functions $\mathbf{f}$ of the form $\mathbf{f}^{(i,j)} = \Delta^{-1} Q_3^{(i)} Q_2^{(j)}$ where, for example,

$$Q_p^{(1)} = \gamma^p, \quad Q_p^{(2)} = \beta^p + \delta^p, \quad Q_p^{(3)} = \beta^p + \gamma^p + \delta^p, \quad Q_p^{(4)} = \beta^p - \gamma^p + \delta^p. \quad (4.38)$$

Some of these possibilities we have already examined: for example, we have confirmed that $\mathbf{f}^{(3,1)}$ generates an $SO(24) \otimes SO(16)$ model, and that $\mathbf{f}^{(1,3)}$, $\mathbf{f}^{(2,3)}$, and $\mathbf{f}^{(3,3)}$ each generate the $SO(24) \otimes E_8$ model. In fact, we have also seen that $\mathbf{f}^{(1,1)}$ leads to the maximally-symmetric $SO(40)$ model (by construction), and $\mathbf{f}^{(4,3)}$ turns out to be a null solution (all the F 's vanish). However, there are eight other distinct modular-invariant functions $\mathcal{Z}$ which can be constructed in this form [eight rather than ten because $\mathbf{f}^{(4,1)}$, $\mathbf{f}^{(4,2)}$, and $\mathbf{f}^{(4,4)}$ each lead to the same $\mathcal{Z}$], and each of these might reasonably correspond to a valid $(1, 4)$ model. For instance, $\mathbf{f}^{(4,1)}$ leads to a modular-invariant function $\mathcal{Z}$ which is the translated version of

$$\mathcal{Z} = \tau_2^{-2} \mathcal{K} (\bar{\beta} \beta^2 - \bar{\gamma} \gamma^2 + \bar{\delta} \delta^2) (\beta^3 - \gamma^3 + \delta^3). \quad (4.39)$$

However, there does not exist a $(1, 2)$ model having this partition function. Our dictionary therefore enables us to construct a $(1, 4)$ model which can be generated for this case.

In a similar manner, we can obtain the corresponding $(1, 4)$ model in $(D = 4)$. In four spacetime dimensions the maximal gauge group is $SO(44)$, and the partition function of the $(1, 4)$ model is (3.13). It is a simple matter to rewrite

$$\begin{aligned} \bar{\beta}^{3/2} \beta^{11/2} + \bar{\gamma}^{3/2} \gamma^{11/2} + \bar{\delta}^{3/2} \delta^{11/2} &= \\ &= \bar{\beta}^{3/2} \beta^{11/2} + \frac{1}{2} (\bar{\gamma}^{3/2} + \bar{\delta}^{3/2}) (\gamma^{11/2} + \delta^{11/2}) \\ &\quad + \frac{1}{2} (\bar{\gamma}^{3/2} - \bar{\delta}^{3/2}) (\gamma^{11/2} - \delta^{11/2}) \end{aligned}$$

whereupon a quick comparison with the solutions of the $K = 8$ dictionary:

$$\begin{aligned} A_{(8)}^1 &\equiv A_8 \stackrel{*}{=} R_1^{K=8} \\ A_{(8)}^2 &\equiv B_8 \stackrel{*}{=} R_2^{K=8} \\ A_{(8)}^3 &\equiv C_8 \stackrel{*}{=} R_3^{K=8} \end{aligned}$$

where the factors from the right-moving $K = 8$ model are

$$\begin{aligned} R_1^{K=2, D=4} &\equiv \frac{1}{4} \Delta^{-1/2} J(\tau) \\ R_2^{K=2, D=4} &\equiv \frac{1}{4} \Delta^{-1/2} J(\tau + \frac{1}{2}) \\ R_3^{K=2, D=4} &\equiv \frac{1}{4} \Delta^{-1/2} J(\tau + \frac{\sqrt{3}}{2}) \end{aligned}$$

Note that it is extremely fortunate that the solution $\mathbf{f}$ satisfies $P_3 S P_3 \mathbf{f} = 0$, for if this solution had not

could not have been built. Once again, we observe that this dictionary is self-consistent, with both sides of (4.41) having modular weight $k = -1$ and mixing identically according the matrices $M_{(8)}$ and $N_{(8)}$ under the $[S]$ - and $[T]$ -transformations respectively. We remark that $(2, 8)$ models can be built as well, simply by taking these expressions $R_i^{K=2, D=4}$ as the valid representative partition-function contributions from self-consistent *left-moving* $K = 2$ sectors.

As in the $K = 4$ case, the J factors in (4.42) arise in the $(1, 2)$ theory from spacetime-related degrees of freedom; the remaining factors, on the other hand, arise from purely internal degrees of freedom. However, unlike the $K = 4$ case, it would have been impossible now to obtain a complete GSO projection with these internal factors, for the analogous combination $\gamma^{3/2} - \beta^{3/2} - \delta^{3/2}$ is non-vanishing. Fortunately, the $K = 8$ theory provides *three* string-function expressions A_8 , B_8 , and C_8 to relate to our three $R^{K=2, D=4}$ factors, so our $K = 8$ dictionary could nevertheless be constructed. It is indeed curious and fortuitous that the $K = 4$ and $K = 8$ theories provide exactly the needed number of independent string-function expressions with which to build supersymmetric fractional superstring partition functions.

5. MODEL TRANSLATION

In the previous section we established our dictionaries, and now we use them to see if they enable us to confirm whether the possible $(1, K)$ models are all valid $(1, K)$ models. Of considerably more interest is to examine those that *do* correspond to models in this space of new $(1, K)$ models and determine their relevant features. This is, in a sense, the opposite of how we use our dictionaries to determine which of the $(1, K)$ models are translated. We will find that not all $(1, 2)$ models are translated, that only those possessing a *maximal* number of spacetime supersymmetries are in correspondence with valid $(1, K)$ models, and that only those with spacetime supersymmetry^[3].

At first glance it may seem that our dictionaries could translate *any* spacetime-supersymmetric $(1, 2)$ model to a valid $(1, K)$ model. For the $K = 4$ case, if our $(1, 2)$ model has a partition function

$$\mathcal{Z}_{(1,2)} = \tau_2^{-2} \Delta^{-1} \overline{\Delta}^{-1/2} \overline{J}(\overline{\tau})$$

where X , Y , and Z are arbitrary holomorphic functions, then the algebraic identity

$$\begin{aligned} \overline{\beta}X + \overline{\gamma}Y + \overline{\delta}Z &= (\overline{\gamma} + \overline{\delta}) \left[\frac{1}{2}(Y + Z) \right] \\ &+ (\overline{\beta} + \overline{\gamma} - \overline{\delta}) \left[\frac{1}{2}X + \frac{1}{4}Y - \frac{1}{4}Z \right] \end{aligned}$$

and the dictionary (4.35) allow us to construct a valid $(1, 4)$ model.

corresponding $(1, 4)$ model:

$$\mathcal{Z}_{(1,4)} = \tau_2^{-2} \Delta^{-1} \left\{ \overline{A}_4 (Y + Z) + \overline{B}_4 [4X + 2(Y - Z)] \right\}. \quad (5.3)$$

This result will be modular-invariant provided (5.1) is modular-invariant. Note that the identity (5.2), which makes the necessary rewriting possible, is simply the statement that the linear combinations $\overline{\gamma} + \overline{\delta}$, $\overline{\beta} + \overline{\gamma} - \overline{\delta}$, and $\overline{\mathcal{J}} \equiv \overline{\gamma} - \overline{\beta} - \overline{\delta}$ span the three-dimensional space spanned by the combinations $\overline{\beta}$, $\overline{\gamma}$, and $\overline{\delta}$ separately, and therefore setting $\overline{\mathcal{J}} = 0$ (the internal GSO projection) always leaves the remaining two left-moving factors F^1 and F^2 to multiply the right-moving factors $\overline{A}_4$ and $\overline{B}_4$ in the $(1, 4)$ partition function. The same argument can be made for the $K = 8$ case as well. It would therefore seem that the spaces of $(1, K)$ models are as large as the spaces of $(1, 2)$ models compactified to the appropriate dimensions $D = 2 + 16/K$, a conclusion which would suggest that the dictionary translation idea does not yield the expected substantial truncation in the sizes of the space of $(1, K)$ models (which are of course *in* their critical dimensions).

Fortunately, this is not the case, for there are two important reasons why valid supersymmetric $(1, 2)$ models may fail to be translatable. First, they may fail to have partition functions of the needed general forms

$$\mathcal{Z} = \tau_2^{-2} \Delta^{-1} \overline{\Delta}^{-1/2} \overline{\mathcal{J}} \left\{ \overline{\beta}^{(10-D)/4} X + \overline{\gamma}^{(10-D)/4} Y + \overline{\delta}^{(10-D)/4} Z \right\}; \quad (5.4)$$

indeed, we will see that the vast majority of known supersymmetric $(1, 2)$ models do not have this form. Second, even though a given model may technically

have a partition function of the form (5.4), its contributions to that partition function may take other forms. We will discuss each of these possibilities.

The first possibility is that a given $(1, 2)$ model may fail to have a partition function of the form (5.4). Spacetime supersymmetry guarantees the presence of ϑ -functions must always appear as in (5.4) for $D = 6$. If we are not assured that our remaining right-moving factors will be $\overline{\beta}^{(10-D)/4}$, $\overline{\gamma}^{(10-D)/4}$, $\overline{\delta}^{(10-D)/4}$, or the combinations from (3.14) and (3.15) we see that we are assured that the factors must contain $10 - D$ powers of ϑ -functions. To obtain (in the $K = 4$ case) expressions such as (5.4) appearing instead. Since the self-consistency of the dictionary demands that the number of remaining ϑ powers be integer, fractional powers are indeed our only other possibility. Supersymmetric $(1, 2)$ models in $D = 6$ will indeed have partition functions. The same problem exists for $D = 4$. Translatability requires that these factors take the form (5.4) yet terms such as $\overline{\beta}\overline{\gamma}^{1/2}$ and $(\overline{\beta}\overline{\gamma}\overline{\delta})^{1/2}$ are also possible occurrences in the space of supersymmetric models. We stand which models do not have such factors will thereby be translatable.

It is clear that in order to obtain the desired

factors –and simultaneously avoid the unwanted ones– our underlying model must have a symmetry relating the internal (*i.e.*, non-spacetime-related) right-moving worldsheet fermions so that they each uniformly produce the same contributions to $\mathcal{Z}$. For example, these $10 - D$ internal fermions must have the same toroidal boundary conditions in all sectors of the model, and be in all respects interchangeable. Such a worldsheet symmetry is not new: if such a symmetry appears amongst the *left-moving* worldsheet fermions, there will be corresponding massless spacetime vector particles transforming in the adjoint representation of this symmetry group. Thus, such a left-moving worldsheet symmetry can be interpreted as a spacetime gauge symmetry. What, however, are the spacetime consequences of such a *right-moving* worldsheet symmetry involving these $10 - D$ worldsheet fermions?

Fortunately, such a worldsheet symmetry is easy to interpret: it is responsible for a multiplicity in the number of spacetime gravitinos, so that the larger the rank of the symmetry group, the larger the multiplicity. In fact, if the rank of this worldsheet symmetry group is $10 - D$ (so that *all* internal right-moving fermions are involved, as is needed for translatability), then the number of gravitinos in the spectrum of the model is N_{max} , where N_{max} is given in (4.1). The analysis needed for proving this assertion is not difficult, but varies greatly with the type of $(1, 2)$ model-construction procedure we employ; in Appendix B we provide a proof using the free-fermion construction of Ref. [10]. We therefore conclude that only models with $N = N_{\text{max}}$ supersymmetry have partition

functions of the form (5.4).^{*}

In the $K = 2$ case ($D = 10$), this of course means that the $N_{\max} = 1$ for $D = 10$, the only models which have $N_{\max} = 1$ (5.4) are indeed those which are supersymmetric. These models, ^[2] and their partition functions are given in Table 1. However, this translatability requirement is much more restrictive than the majority of supersymmetric $(1, 2)$ models in $D = 10$ are not SUSY. For example, it has been found ^[12] that there are only 10 of $D = 4$ free-fermion models, there exist fewer than 100 of $D = 6$ free-fermion models, this is to be compared against over 32 000 models in $D = 10$. This is a virtually limitless supply with no spacetime supersymmetry. Therefore, that the number of translatable $(1, 2)$ models is small compared to the number of $(1, 8)$ models— is severely restrictive. A similar restriction exists for the $K = 4$ case.

As mentioned earlier, there is also a second way in which the partition function may fail to be translatable, even if it does technically satisfy the axioms of the algebraic form (5.4). Let us consider the second example, the spacetime partition function of a model from the *spacetime* Jacobson paper, respectively. Recall that in the spacetime Jacobson paper, the partition function contribution from spacetime bosons (the Neveu-Schwarz sector) is given by

★ This statement assumes that we have canonical time bosonic and fermionic states in contact with the system as will be discussed below.

holomorphic factors in the partition function represent purely internal degrees of freedom. Let us therefore separate these two pieces, and write the general bosonic and fermionic partition functions from an arbitrary model as follows:

$$\begin{aligned}\mathcal{Z}_b &= \tau_2^k \Delta^{-1} \overline{\Delta}^{-1/2} \left\{ (\overline{\gamma} - \overline{\delta}) V + W_b \right\}, \\ \mathcal{Z}_f &= \tau_2^k \Delta^{-1} \overline{\Delta}^{-1/2} \left\{ \overline{\beta} V + W_f \right\}.\end{aligned}\tag{5.5}$$

Here W_b is that part of $\mathcal{Z}_b$ whose spacetime anti-holomorphic factor is not proportional to $(\overline{\gamma} - \overline{\delta})$. It is clear that the total partition function $\mathcal{Z} \equiv \mathcal{Z}_b - \mathcal{Z}_f$ will be of the algebraic form (5.4) if $W_b = W_f \equiv W$ where W is arbitrary and if V is of the form of the term in braces in (5.4). However, translatability additionally requires that W vanish, regardless of its form. The reason for this is quite simple. If $W \neq 0$, then allowing W_b and W_f to cancel in the difference $\mathcal{Z}_b - \mathcal{Z}_f$ amounts to cancelling spacetime bosonic and fermionic states and is therefore inconsistent with our retention of the spacetime Jacobi factor $\overline{J}$ in (5.4); indeed, writing $W_b - W_f = 0$ is tantamount to simply writing $\overline{J} = 0$ in (5.4). Another way of seeing this is to realize that while W_b and W_f might be algebraically equal, the *spacetime* factor in W_b must include combinations of $\overline{\gamma}$ and $\overline{\delta}$ only (since it comes from spacetime bosonic degrees of freedom), while the spacetime factor in W_f must be simply $\overline{\beta}$. Such expressions W_b and W_f should therefore not be cancelled, so translatability requires that they not appear at all. Note that this additional constraint $W = 0$ is not vacuous, for there exist many models for which the total $\mathcal{Z}$ is of the algebraic form (5.4) but for which $W \neq 0$: these are models with partition functions of the form

$\mathcal{Z} + W - W$. It turns out, however, that any $(1, K)$ model is guaranteed to have $W = 0$, so the $N = N_{\max}$ constraint ensures model translatability. (In fact, as noted above, even for $N < N_{\max}$ does the $N = N_{\max}$ constraint cease to be *overconstraining* again that $N = N_{\max}$ SUSY is the $(1, 2)$ physical model.)

One might argue that this $N = N_{\max}$ translatability constraint is artificial: our dictionary was constructed by requiring that the $(1, K)$ model with the maximally symmetric sector be the only one possible, since this latter model always has an $N = N_{\max}$ sector. However, the $N = N_{\max}$ constraint was thereby “encoded” in the dictionary from the beginning. Indeed, one might claim that other dictionaries would be possible for $N < N_{\max}$ cases simply by comparing, for example, the $(1, K)$ model with a maximally symmetric sector to a $(1, 2)$ model. However, such approaches ultimately fail to yield a dictionary. On a mathematical level, this occurs because it is not possible to write ϑ -function expressions which could appear on the right-hand side of the dictionary: such expressions would have to be non-holomorphic, and would not also be closed under $[S]$, and the $N = N_{\max}$ sector would not be the only ones possible. On a physical level, this is also true. It follows. Unlike the $(1, 2)$ models in $D < 10$, the $(1, K)$ models in critical dimensions, and therefore they lack “internal” degrees of freedom. Thus, we expect that any dictionary mapping a $(1, K)$ sector to a $K > 2$ right-moving sector should

tra degrees of freedom in the $K = 2$ sector in order to introduce twists or (super)symmetry-breaking; rather, we expect these degrees of freedom to be “frozen out”, handled jointly as though they were one block. This is precisely how they appear in the dictionaries (4.35) and (4.41), and is the root of the previously encountered indistinguishability of (or symmetry relating) the internal right-moving fermions. Thus, it is indeed sensible that our $(1, K)$ models are the analogues of the $N = N_{\max}$ $(1, 2)$ models, for both are the unique models in which no internal right-moving degrees of freedom are available for (super)symmetry breakings. Note that this argument does not tell us the degree of supersymmetry for the $(1, K)$ models themselves. However, both an examination of the individual bosonic and fermionic degrees of freedom and an overall counting of states indicate that the $(1, K)$ strings have $N = 1$ supersymmetry [as distinguished from the (K, K) strings, which have $N = 2$ supersymmetry^[31]].

Finally, we should point out that it is also possible to build fractional superstring models which are *not* spacetime-supersymmetric. There are primarily two ways in which this can be done. First, it is possible to construct string-function expressions which are similar to A_K , B_K , and C_K but which do *not* vanish; partition functions built with these expressions would then correspond only to *non*-supersymmetric models, and analogous dictionaries could be constructed (in the manner presented in Sect. IV) guaranteeing that such self-consistent $(1, K)$ models actually exist. Constructing such expressions is not difficult: of all the requirements listed after (3.27), we need only eliminate the tachyon-free constraint $a_n = 0$ for all $n < 0$. Note that removal of this require-

ment need not introduce spacetime tachyons in the $(1, K)$ model, for in general a partition function

$$\mathcal{Z} = \tau_2^k \sum_{m,n} a_{mn} \bar{q}^m$$

will have physical tachyons only if a *diagonal* element $a_{nn} \neq 0$ for $n < 0$. Indeed, it is easy to choose holomorphic functions a_{mn} in such a way that $a_{nn} = 0$ for all $n < 0$.

A second way to build non-supersymmetric models is to start with the supersymmetric expressions A_K , B_K , and C_K and then separate them into their spacetime bosonic and fermionic parts in a way that $A_K = A_K^{(b)} - A_K^{(f)}$, *etc.* For the $K = 2$ case, it is clear how to do this: the term β reappears in the Ramond (*i.e.*, spacetime-fermionic) sectors in the $\gamma - \delta$ term which arises only from a Neveu-Schwarz sector. For the $K > 2$ “para-Jacobi” identities $A_K = A_K^{(b)} - A_K^{(f)}$, the situation is more complicated. A first approach would be to develop in Sect. IV to relate our desired $K > 2$ models to the known $K = 2$ pieces: we would simply split the para-Jacobi factor J which appears in the $K = 2$ models into two parts, attempt to construct, for example, the $K = 4$ spacetime-fermionic $B_4^{(b,f)}$ and $B_4^{(b,f)}$ which would be consistent with the

$$\begin{aligned}
A_4^{(b)} &\stackrel{*}{=} 4k \Delta^{-1/2} (\gamma - \delta)_{\text{st}} (\gamma + \delta)_{\text{int}} \\
A_4^{(f)} &\stackrel{*}{=} 4k \Delta^{-1/2} (\beta)_{\text{st}} (\gamma + \delta)_{\text{int}} \\
B_4^{(b)} &\stackrel{*}{=} k \Delta^{-1/2} (\gamma - \delta)_{\text{st}} (\beta + \gamma - \delta)_{\text{int}} \\
B_4^{(f)} &\stackrel{*}{=} k \Delta^{-1/2} (\beta)_{\text{st}} (\beta + \gamma - \delta)_{\text{int}}
\end{aligned} \tag{5.7}$$

where we have explicitly indicated with subscripts the separate $K = 1$ spacetime and internal factors. However, it is easy to see that this method is not appropriate, for we do not expect the cancellations occurring in the full expressions A_K and B_K to mirror the relatively simple cancellation occurring in the $K = 2$ case. Indeed, it is easy to show that no string functions for the left sides of (5.7) can be found which transform under $[S]$ and $[T]$ as do the right sides. Instead, one can determine the separate bosonic and fermionic contributions to the expressions A_K , B_K and C_K by demanding that these individual contributions each have q -expansions $\sum_n a_n q^n$ in which all coefficients a_n are non-negative, and have relatively simple closure relations under $[S]$. This approach, in fact, proves successful, and yields results consistent with our interpretation [discussed after (3.31)] that the terms $(c_0^0)^{D-3}(c_0^2)$ are bosonic and the terms $(c_{K/2}^{K/2})^{D-2}$ are fermionic.^[11] Thus, the construction of *non*-supersymmetric $(1, K)$ models is in principle no more difficult than that of the supersymmetric models we have already considered, and we expect our dictionary techniques to generalize to these cases as well [though of course not yielding dictionaries similar to (5.7)].

To summarize the results of this and the previous section, then, we have succeeded in developing a method by which the partition functions of valid supersymmetric $(1, K)$ models can be generated and their gauge groups identi-

fied. This was achieved by establishing a correspondence between the $(1, 2)$ models in $D = 2 + 16/K$ dimension and the $(1, K)$ models in $D = 2 + 16/K$ dimension [Eqs. (4.35) and (4.41)] enabling one to “translate” the results of the well-understood $K = 2$ sector and the less-understood $K > 2$ sector. We found that only $(1, 2)$ models are translatable, and we were thereby able to establish the validity of the $(1, K)$ models.

6. FURTHER DISCUSSION AND REMARKS

In this concluding section we discuss two different extensions of our results: these are the questions of obtaining chiral fermions and achieving Lorentz invariance. We begin by investigating how chiral fermions – or more generally, chiral supermultiplets – might arise in the fractional superstring models we have been investigating. We discuss several different methods for obtaining such multiplets, one of which involves formulating or interpreting these models in dimensions less than their critical dimensions. While this might seem to spoil the original attraction of the fractional superstring approach, we find instead that such a compactification is not at all arbitrary (as it is for the traditional superstrings), but rather is required in order to achieve a self-consistent Lorentz-invariant interpretation. Indeed, we find that requiring Lorentz invariance seems to specify a “natural” dimension in which the theory must be formulated, thereby (in a unique manner) simultaneously offering a possible solution to the chiral fermion problem. We emphasize that such a “forced” compactification appears to be a feature wholly new to fractional superstrings. Furthermore, we find that the “natural” dimension for the $K = 4$ string appears to be $D = 4$, rendering the $K = 4$ fractional superstring the most likely candidate for achieving chiral particle representations in four-dimensional spacetime while maintaining Lorentz invariance.

We begin by investigating how we might obtain chiral massless spacetime supermultiplets (*i.e.*, supermultiplets which transform in a complex representation of the gauge group) in our fractional superstrings. Recall from Sect. III that

the massless particles appear only in the A_K supermultiplets we have examined; indeed, for $K \geq 2$, the $(c_{K/2}^0)^{D_c-2}$ term contains the contributions of the right-moving component particles A^μ while the $(c_{K/2}^{K/2})^{D_c-2}$ term contains the ψ^α (where α is a spacetime Lorentz spinor index). In this theory in its critical dimension, the massless spectrum is in general

$$(\psi^\alpha, A^\mu)_R \otimes \left\{ X^\nu \oplus \phi^{(h_1)} \oplus \dots \right\}$$

here $(\psi^\alpha, A^\mu)_R$ is the fermion/vector supermultiplet of the left-moving excitations we have indicated in (6.1). The states: X^ν denotes the Lorentz-vector (gauge-singlet) the worldsheet left-moving component of the supermultiplet; the $\phi^{(h_i)}$ denote the various remaining Lorentz representations (with highest weights h_i) of the right-moving. Within (6.1), the combination

$$(\psi^\alpha, A^\mu)_R \otimes (X^\nu)$$

forms the usual $N = 1$ supergravity multiplet consisting of $g^{\mu\nu}$, spin-1 antisymmetric tensor field $B^{\mu\nu}$, spin-1 $\lambda^{\nu\alpha}$, and spin-1/2 fermion λ^α . All of these states are massless. The only other combination within (6.1) allows

is

$$(\psi^\alpha, A^\mu)_R \otimes (\phi^{(h_{\text{adj}})})_L \quad (6.3)$$

where these left-moving states fill out the adjoint representation of the gauge group. This yields, of course, a spacetime supermultiplet consisting of spin-1 gauge bosons A^μ_{adj} and their spin-1/2 fermionic superpartners ψ^α_{adj} . The crucial point, however, is that this supermultiplet structure therefore does not permit spacetime fermions which transform in any other (*e.g.*, complex) representations of the gauge group.

In the usual superstring phenomenology, a chiral supermultiplet can be achieved by introducing an additional right-moving supermultiplet of the form $(\phi_i, \psi^\alpha)_R$ where the ϕ_i , $i = 1, \dots, D-2$, are a collection of Lorentz scalar fields. We then obtain, as in (6.1), the additional states

$$(\phi_i, \psi^\alpha)_R \otimes \left\{ X^\nu \oplus \phi^{(h_1)} \oplus \phi^{(h_2)} \oplus \dots \right\}_L. \quad (6.4)$$

The combination

$$(\phi_i, \psi^\alpha)_R \otimes (X^\nu)_L \quad (6.5)$$

yields additional spin-1 vector bosons $\tilde{A}_i^\nu$, spin-1/2 fermions ψ^α , and spin-3/2 gravitinos $\lambda^{\nu\alpha}$ (thereby producing an $N > 1$ SUSY unless these extra gravitinos are GSO-projected out of the spectrum). However, since (6.4) is built with the right-moving supermultiplet $(\phi_i, \psi^\alpha)_R$, the level-matching constraints now also

allow the additional combinations

$$(\phi_i, \psi^\alpha)_R \otimes (\phi^{(h_{\text{adj}})})_L$$

where the gauge-group representations are *not* the adjoint. These combinations (6.6) can yield chiral spacetime fermions with the highest weights of complex representations and thus contain chiral massless spacetime fermions transforming in the fundamental representation.

How might these additional supermultiplets arise in superstring theories? At the partition function level, new terms of the form $(c_0^0)^{D_c-3}(c_0^2)$ to be interpreted as excitations of massless scalar fields. Such (ϕ_i, ψ^α) multiplets are introduced simply by multiplying the overall $K = 4$ by an appropriate integer (so as to maintain modular invariance). In the actual particle spectrum, however, it is not clear how they arise. Extra fermions are not difficult to find, but extra degrees of freedom while the Dirac γ -matrix algebra, and the level-matching condition, reduces this to $2^{D/2-1}$ degrees of freedom. The other degrees of freedom remaining for chiral fermions and scalar fields, on the other hand, are more difficult to accommodate. As follows. For $K > 2$ there also exist extended representations which have the same central charge as our usual representations. More than one ϕ_0^1 field. At first sight, these extended

Lorentz indices μ associated with them. However, recall that in the parafermion theories there are additional parafermion fields whose characters have not appeared in the partition functions: these are the half-integer spin fields. It may therefore be possible to interpret these extra ϕ_0^1 fields as Lorentz scalar *composites* of the half-integer spin fields – *e.g.*, the composite $(\phi_{1/2}^{1/2})^\mu (\phi_{-1/2}^{1/2})_\mu$ may contain the additional ϕ_0^1 field in the extended parafermion theory. Forming these ϕ_0^1 fields as composites may allow them to be interpreted as Lorentz scalars, much as the scalar composite $:\epsilon^\mu\epsilon_\mu:$ appears as one of the descendants of ϵ^μ [as in (2.10)]. These scalars would then supply the scalar fields needed for chiral supermultiplets. Of course, a detailed analysis is necessary to see if these scenarios are possible.

A more widely-known and established method of generating supermultiplets (ϕ_i, ψ^α) in string theory is through spacetime compactification. Since the $K = 8$ string is already in $D_c = 4$, let us focus on the $(1, 4)$ heterotic string with $D_c = 6$. For this string, we can choose to compactify two space dimensions:

$$(\psi^\alpha, A^\mu) \longrightarrow (\psi^\alpha, A^\mu) \oplus (\phi_i, \psi^\alpha); \quad (6.7)$$

here the A^μ , $\mu = 0, 1, \dots, 5$ in the $D = 6$ theory breaks down to A^μ , $\mu = 0, 1, 2, 3$ in $D = 4$, with the remaining fields reidentified as internal scalars: $A_4 = \phi_1$ and $A_5 = \phi_2$. Similarly, the four-component Weyl fermion in $D = 6$ splits into two two-component fermions in $D = 4$. Therefore, the degree-of-freedom count corresponding to (6.7) is

$$(4 + 4) \longrightarrow (2 + 2) + (2 + 2). \quad (6.8)$$

In the usual $(1, 2)$ string compactified to four dimensions, it is precisely this mechanism which breaks the ten-dimensional Lorentz symmetry to the four-dimensional Lorentz symmetry. The separation of the $(\phi_i, \psi^\alpha)_R$ can then couple to different left-moving representations of chiral fermions in the fundamental representation of the gauge group.

There is, however, a crucial difference between the usual $(1, 2)$ heterotic string and the compactified string in four spacetime dimensions. In the usual superstring theory, one arbitrarily chooses the resulting dimensionality of spacetime to be the $(1, 2)$ theory, and there are neither doubts nor why four dimensions is chosen. For the $(1, 4)$ heterotic string, that even though the critical dimension is $D_c = 10$, it indicates that there does not exist a six-dimensional interpretation consistent with Lorentz invariance and supersymmetry. Indeed, we will see that the largest number of dimensions for which a consistent interpretation exists is $D = 4$, the *critical dimension*. Hence, the $(1, 4)$ string appears unusual to (or interpretation in) four spacetime dimensions because of its intrinsic structure of the theory itself.

Let us see how this comes about. Recall that the character for $[\phi_m^j]$. Also recall that the parafermion ϕ_m^j has the m -quantum number is additive (modulo k)

ber is not. As we have already observed, the $(c_0^0)^3(c_0^2)$ term in A_4 represents the contributions from spacetime bosons, and the $(c_2^2)^4$ term in A_4 represents the contributions from spacetime fermions. Hence, as is argued in Ref. [11], terms with $D - 2$ factors of $c_{K/2}^\ell$ (with arbitrary values of ℓ) correspond to spacetime fermions, while terms of the forms $(c_0^\ell)^{D-2}$ (with arbitrary values of ℓ) correspond to spacetime bosons. Such identifications arise essentially from recognizing the worldsheet vacuum state corresponding to $(\phi_0^0)^{D-2}$ as yielding spacetime bosons [it is the analogue of the $K = 2$ Neveu-Schwarz vacuum $(\phi_0^0)^8$], and recognizing the worldsheet vacuum state corresponding to $(\phi_{\pm K/4}^{K/4})^{D-2}$ as yielding spacetime fermions [it is the analogue of the $K = 2$ Ramond vacuum $(\phi_{\pm 1/2}^{1/2})^8$]. Note that this identification is in fact consistent at all mass levels, since excitations upon these respective vacua occur only through the $\epsilon = \phi_0^1$ field, and the parafermion fusion rules show that applications of ϕ_0^1 cannot change the m -quantum numbers of these respective vacua. Furthermore, since the m -quantum number is additive modulo $K/2$, this identification indeed reproduces the desired Lorentz fusion rules $B \otimes B = B$, $B \otimes F = F$, and $F \otimes F = B$, where B and F represent spacetime bosonic and fermionic fields. Thus, with this interpretation in $D = D_c$ spacetime dimensions, we see that it is straightforward to determine the spacetime statistics of the particles contributing to each term in A_4 and A_8 .

However, we now note that in B_4 there exists a term of the form $(c_2^2)^2(c_0^2)^2$ which has only massive states [*i.e.*, in a q -expansion of this term we find $q^h(1 + \dots)$ where $h > 0$]. Although this term may have more than one interpre-

tation, the elementary particles giving rise to the term arise through ϵ -excitations from either the six-dimensional vacuum state. It is, therefore, natural to interpret the term from either:

1. fermions in four spacetime dimensions [the term arises from $(\phi_0^0)^4$ with additional internal quantum numbers],
2. bosons in four spacetime dimensions [the term arises from $(\phi_0^0)^4$ with additional internal quantum numbers].

How might such internal factors arise? Let us consider the case of heterotic superstring theory, where in $D_c = 10$ spacetime dimensions the spinor (corresponding to spacetime fermions) always carries an internal index. In the function the factor $-(c_1^1)^8 = -\vartheta_2^4/(16\Delta^{1/2})$. This is missing in A_2 in Eq. (3.29); the presence of the term is a reflection of the violation of the underlying rotational invariance of the theory. However, compactifying a model to a dimension $D < D_c$ is required; indeed, we have seen in Sect. III that in $D < D_c$ flat spacetime dimensions, we obtain terms in the expansion with string-function factors of the form $c^{D-2}c^L$ arising from spacetime-associated degrees of freedom (where L is the number of spacetime Lorentz indices) and where the term carries additional internal degrees of freedom (*i.e.*, worldsheet indices). In the present case, upon compactification

the form $(c_{K/2}^\ell)^{D-2}(c_0^{\ell'})^{D_c-D}$ where the second factor represents contributions from spacetime bosons (and is the $K > 2$ analogue of the contributions from the usual Neveu-Schwarz sectors of the theory). Remarkably, this is precisely what occurs in B_4 . Apparently, six-dimensional Lorentz invariance for the $K = 4$ string appears to be explicitly broken.

Note that – at tree-level – the particles in the A_4 sector couple only to other fields in the A_4 sector; in particular, the tree-level scatterings of massless particles do not involve the B_4 sector. At the string *loop* level, however, fields in the B_4 sector couple to the massless particles in the A_4 sector, so we see that it is the string’s own quantum effects which render the theory in its critical dimension inconsistent with Lorentz invariance. Thus, in this sense, it is quantum mechanics and the requirement of Lorentz invariance which together necessitate or *induce* the compactification from the critical dimension (six) to the natural dimension (four). This mechanism is quite remarkable, and has no analogue in the traditional $K = 1, 2$ string theories (in which the critical dimensions *are* the natural dimensions). In particular, only for the $K > 2$ theories do such independent B -sectors appear; for $K = 2$ there is only one possible supersymmetric A sector, corresponding to the one supersymmetry identity (Jacobi identity) for $K = 2$. It is the presence of a unique B_K sector for $K > 2$, itself necessitated by modular invariance and closure under $[S]$, which induces this compactification.

The same argument applies to the $K = 8$ string (for which the critical dimension is four but the natural dimension is three), but with one additional

feature. As before, the A_8 expression [Eq. (3.3)] is consistent with a $D = D_c = 4$ spacetime boson/fermion counting; we can quickly see that the B_8 expression is *not* consistent; rather, the form of this expression dictates a different interpretation. Now, the *third* expression C_8 involves only terms of the type $c_{K/4}^\ell = c_2^\ell$. (Recall that the only C_8 expression consistent with three-dimensional Lorentz invariance is in fact Lorentz invariant in any number of dimensions.) The above interpretation indicates that C_8 must correspond to a spacetime Lorentz spins $1/4$ and $3/4$. However, spin $1/4$ fields (i.e., anyons) are consistent with Lorentz invariance in three spacetime dimensions, though not in four. The way in which self-consistency selects this dimensionality for $K > 2$ strings seems to select naturally their proper critical dimensions are the dimensions in which consistent Lorentz invariance is achieved.

Note that if such speculations are indeed correct, the models we have constructed in this paper have a very different spectra, for when we interpret them in four dimensions, respectively these string theories contain worldsheet bosons compactified. However, we expect the “dictionary” between the models in this paper to be easily generalizable to the other cases as well.

In summary, then, although there may well exist other possible interpretations of these partition functions, we find the above speculations both tantalizing and intriguing. It is indeed fortuitous that the solution to what might have seemed a problem (*i.e.*, finding an interpretation consistent with Lorentz invariance) also simultaneously provides a possible solution to the more phenomenological problem of obtaining models with chiral fermions. It is also strongly compelling that the quantum structure of the theory itself seems to dictate this solution. These are issues clearly worth investigating.

ACKNOWLEDGE

We are pleased to thank S.-W. Chung, J. C. S. Lam, E. Lyman, R. Myers, and especially P. ... sions. K.R.D. also thanks the High-Energy The ... for its hospitality while portions of this work v ... supported in part by the U.S. National Scienc ... ence and Engineering Research Council of Ca ... Québec.

APPENDIX A

In this appendix we introduce the string functions (or parafermion characters) c_n^ℓ , collecting together their definitions, modular transformation properties, special cases, and identities. The results quoted here are due mostly to Kač and Peterson.^[9]

The string functions are essentially the characters Z_{2m}^{2j} of the $\mathbf{Z}_K$ parafermion fields ϕ_m^j :

$$Z_n^\ell = \eta c_n^\ell \quad (\text{A.1})$$

where $\ell \equiv 2j$, $n \equiv 2m$, and where η is the Dedekind η -function:

$$\eta(\tau) \equiv q^{1/24} \prod_{n=1}^{\infty} (1 - q^n) = \sum_{n=-\infty}^{\infty} (-1)^n q^{3(n-1/6)^2/2} \quad (\text{A.2})$$

with $q \equiv e^{2\pi i \tau}$. Since the $\mathbf{Z}_K$ parafermion theory can be represented as the coset theory $SU(2)_K/U(1)$, the string functions can therefore be obtained by expanding the full $SU(2)_K$ characters $\chi_\ell(\tau, z)$ in a basis of $U(1)$ characters $\Theta_{n,K}(\tau, z)/\eta$:

$$\chi_\ell(\tau, z) = \sum_{n=-\ell}^{2K-\ell-1} Z_n^\ell(\tau) \frac{\Theta_{n,K}(\tau, z)}{\eta} = \sum_{n=-\ell}^{2K-\ell-1} c_n^\ell(\tau) \Theta_{n,K}(\tau, z). \quad (\text{A.3})$$

Here $\Theta_{n,K}(\tau, z)/\eta$ are the characters of a $U(1)$ boson compactified on a radius $\sqrt{K}$. Since the $SU(2)_K$ characters and the $\Theta_{n,K}$ functions are well-known, explicit expressions for the string functions can be extracted. Eq. (A.3), then, can be taken as a definition of the string functions c_n^ℓ .

Expressions for the string functions c_n^ℓ were given by Peterson;^[9] for our purposes, however, a useful expression is due to Distler and Qiu:^[15]

$$c_n^\ell(\tau) \equiv q^{h_n^\ell + [4(K+2)]^{-1}} \eta^{-3} \sum_{r,s=0}^{\infty} (-1)^{r+s} q^{r(j+m) + s(j-m)} - q^{K+1-2j+r}$$

where $\ell - n \in 2\mathbf{Z}$ and where the highest weight h_n^ℓ is

$$h_n^\ell \equiv \frac{\ell(\ell+2)}{4(K+2)} - \frac{n^2}{4K}$$

The string functions have the symmetries

$$c_n^\ell = c_{K-n}^{K-\ell} = c_{-n}^\ell =$$

as a consequence of which for any K we are free to choose the string functions c_n^ℓ where $0 \leq \ell \leq K$ and $0 \leq n \leq n_{\max}$ and $\ell - 2$ otherwise].

The string functions are closed under the modular transformation T . In fact, under $T: \tau \rightarrow \tau + 1$ they transform as

$$c_n^\ell(\tau + 1) = \exp\{2\pi i s_n^\ell\} c_n^\ell(\tau)$$

where the phase s_n^ℓ is given by

$$s_n^\ell \equiv h_n^\ell - \frac{1}{24} c_0 = h_n^\ell - \frac{1}{24} c_0$$

Here h_n^ℓ is defined in (A.5), and c_0 is the central charge.

theory [given in (2.13)]. Under $S : \tau \rightarrow -1/\tau$, the level- K string functions mix among themselves:^[9]

$$c_n^\ell(-1/\tau) = \frac{1}{\sqrt{-i\tau}} \frac{1}{\sqrt{K(K+2)}} \sum_{\ell'=0}^K \sum_{\substack{n'=-K+1 \\ \ell'-n' \in 2\mathbf{Z}}}^K b(\ell, n, \ell', n') c_{n'}^{\ell'} \quad (\text{A.9})$$

where the first square root indicates the branch with non-negative real part and where the mixing coefficients $b(\ell, n, \ell', n')$ are

$$b(\ell, n, \ell', n') \equiv \exp\left\{\frac{i\pi n n'}{K}\right\} \sin\left\{\frac{\pi(\ell+1)(\ell'+1)}{K+2}\right\}. \quad (\text{A.10})$$

From (A.7) and (A.9) it follows that the string functions have modular weight $k = -\frac{1}{2}$.

Note that in the special $K = 1$ and $K = 2$ cases, the string functions can be expressed in terms of the Dedekind η -function (A.2) and the more familiar Jacobi ϑ -functions; these relations are given in (3.18) and (3.19). In particular, these ϑ -functions are

$$\begin{aligned} \vartheta_2(\tau) &\equiv 2 q^{1/8} \prod_{n=1}^{\infty} (1 + q^n)^2 (1 - q^n) \\ \vartheta_3(\tau) &\equiv \prod_{n=1}^{\infty} (1 + q^{n-1/2})^2 (1 - q^n) \\ \vartheta_4(\tau) &\equiv \prod_{n=1}^{\infty} (1 - q^{n-1/2})^2 (1 - q^n), \end{aligned} \quad (\text{A.11})$$

and under the $[S]$ and $[T]$ transformations [where the bracket indicates the

stroke operator, defined in (3.10)], we find

$$\begin{aligned} \vartheta_2[S] &= \exp(7\pi i/4) \vartheta_4, & \vartheta_2[T] &= \exp(7\pi i/4) \vartheta_4, \\ \vartheta_3[S] &= \exp(7\pi i/4) \vartheta_3, & \vartheta_3[T] &= \exp(7\pi i/4) \vartheta_3, \\ \vartheta_4[S] &= \exp(7\pi i/4) \vartheta_2, & \vartheta_4[T] &= \exp(7\pi i/4) \vartheta_2, \\ \eta[S] &= \exp(7\pi i/4) \eta, & \eta[T] &= \exp(7\pi i/4) \eta. \end{aligned}$$

Note that (A.12) is indeed consistent with the (A.7) and (A.9) for $K = 2$; in particular, under $c_0^0 + c_0^2$ transforms as an eigenfunction while c_0^0 other.

The S -transformation formula (A.9) assumes for the $K = 4$ case. In this case there are several under $[S]$ they can be block-diagonalized as follows

$$\begin{pmatrix} c_0^0 + c_0^4 \\ c_0^2 \\ c_2^4 \\ c_2^2 \end{pmatrix} [S] = e^{i\pi/4} \mathbf{R}^+ \begin{pmatrix} c_0^0 + c_0^4 \\ c_0^2 \\ c_2^4 \\ c_2^2 \end{pmatrix}$$

and

$$\begin{pmatrix} c_0^0 - c_0^4 \\ c_1^1 \\ c_1^3 \end{pmatrix} [S] = e^{i\pi/4} \mathbf{R}^- \begin{pmatrix} c_0^0 - c_0^4 \\ c_1^1 \\ c_1^3 \end{pmatrix}$$

where

$$\mathbf{R}^+ \equiv \frac{1}{\sqrt{24}} \begin{pmatrix} 2 & 4 & 4 & 4 \\ 2 & -2 & 4 & -2 \\ 1 & 2 & -2 & -2 \\ 2 & -2 & -4 & 2 \end{pmatrix}, \quad \mathbf{R}^- \equiv \frac{1}{\sqrt{24}} \begin{pmatrix} 2 & 4 & 4 & 4 \\ 2 & -2 & 4 & -2 \\ 1 & 2 & -2 & -2 \\ 2 & -2 & -4 & 2 \end{pmatrix}$$

Note that $(\mathbf{R}^+)^2 = (\mathbf{R}^-)^2 = \mathbf{1}$, as required.

This mixing pattern seen for the $K = 2$ and $K = 4$ cases exists for other higher even values of K as well. For any such K , we can define the linear combinations $d_n^{\ell\pm} \equiv c_n^\ell \pm c_n^{K-\ell}$ when (ℓ, n) are even; note that these d -functions will also be eigenfunctions of $[T]$ if $K \in 4\mathbf{Z}$ because in these cases $s_n^\ell = s_n^{K-\ell} \pmod{1}$. It then follows from (A.9) that the d^+ -functions mix exclusively among themselves under S [as in (A.13)], and that the d^- -functions mix exclusively with themselves and with the odd (ℓ, n) string functions [as in (A.14)]. It is for this reason that (for $K > 2$) it is possible to construct modular-invariant expressions involving the string functions in such a manner that the odd string functions do not appear. Indeed, one can check that in all the string-function expressions which have appeared for the fractional superstring (such as A_K , B_K , and C_8), only the d^+ combinations have played a role.

APPENDIX

In this appendix we demonstrate, using the construction of Ref. [10], that any $(1, 2)$ model with a partition function of the form

$$\mathcal{Z} = \tau_2^{-2} \Delta^{-1} \overline{\Delta}^{-1/2} \overline{J} \left\{ \overline{\beta}^{(10-D)/4} X + \overline{\gamma}^{(10-D)/4} Y \right\}$$

has an $N = N_{\max}$ spacetime supersymmetry. We assume the reader to be familiar with the model described in Ref. [10], and we use the same notation.

It is clear that a model with partition function $\mathcal{Z}$ is modular invariant; therefore, as discussed in Ref. [10], its $\mathbf{W}_i$ vectors must include the two vectors

$$\begin{aligned} \mathbf{W}_0 &\equiv [(\tfrac{1}{2})^{12+2k} \mid (\tfrac{1}{2})^{4+2k}] \\ \text{and } \mathbf{W}_1 &\equiv [(0)^{-k} (0 \tfrac{1}{2})^{4+2k} \mid (0 \tfrac{1}{2})^{4+2k}] \end{aligned}$$

where k , the modular weight, equals $1 - D/2$. The first two vectors indicate the boundary conditions of the left-moving fermions, and we see that the four complex boundary conditions in $\mathbf{W}_1$ (*i.e.*, the four whose components are the ones which together contribute the spacetime supersymmetry) and the remaining $2(4+k) = 10 - D$ right-moving complex boundary conditions (those whose components in $\mathbf{W}_1$ equal $\tfrac{1}{2}$) are the remaining factors of $\beta^{(10-D)/4}$, $\gamma^{(10-D)/4}$, and $\overline{\beta}^{(10-D)/4}$.

In order for our model to have a partition function of the form (B.1), these remaining $(10 - D)$ worldsheet fermions must have the *same* toroidal boundary conditions in all sectors of the model which contribute to the partition function. In particular, this must be true *separately* for spacetime bosonic and fermionic sectors (*i.e.*, those which have $\alpha s = \frac{1}{2}, 0$ respectively in the notation of Ref. [10]). This can occur only if every other spin-structure generating vector $\mathbf{W}_i$ has equal components for these $10 - D$ fermions (thereby giving rise to the maximal right-moving “gauge” symmetry discussed in Sect. V). Thus, any other spin-structure vectors $\mathbf{W}_i$ in the generating set must have a right-moving component of the form

$$\mathbf{W}_i^R = [(0)^{-k} (X)^{4+k}] \quad (\text{B.3})$$

where each factor X is independently either (000) or $(0\frac{1}{2}\frac{1}{2})$. In determining (B.3) we have had to satisfy the “triplet” constraint; we have also made use of our freedom to choose (without loss of generality) vectors which have their first components vanishing.

Let us now consider the constraints which must be satisfied by the gravitino states. These states all arise in the $\mathbf{W}_1$ sector, and have the charge vectors

$$\mathbf{Q}_{\text{grav}} \equiv \mathbf{N}_{\mathbf{W}_1} + \mathbf{W}_1 - \mathbf{W}_0 = [(\pm\frac{1}{2})^{-k} (\pm\frac{1}{2}00)^{4+k} \mid (0)^{24+k}] \quad (\text{B.4})$$

where in principle all $\pm\frac{1}{2}$ combinations are allowed. The constraint due to the $\mathbf{W}_0$ vector, however, immediately projects out half of these states, allowing only those with charge vectors in which the product of non-zero components is positive. This by itself leaves us with eight distinct allowed states, and these states

are precisely those which combine to form the $\mathbf{W}_1$ vector. Thus, our model will contain *fewer* gravitinos if and only if the generating set produce constraint equations removed from the physical spectrum. The $\mathbf{W}_1$ -vectors are, however, and since any additional $\mathbf{W}$ -vectors in the generating set have components of the forms (B.3), they each give rise to a constraint of the form

$$0 = f(k_{ij})$$

where $f(k_{ij})$ is a function of the projection coefficients. Note, in particular, that (B.5) does not involve k_{ij} for $i, j = 1, 2, 3$. We see that either *all* or *none* of the N_{max} gravitinos remain in the physical spectrum: all gravitinos remain if the projection coefficients are all zero in each case $f(k_{ij}) = 0$, and none remain otherwise. Thus, our model must have at least an $N = 1$ spacetime supersymmetry in order to have a partition function of the form (B.1) with $N = N_{\text{max}}$ spacetime supersymmetry.

REFERENCES

1. See, *e.g.*, M.B. Green, J.H. Schwarz and E. Witten, *Superstring Theory*, Vols. 1 and 2 (Cambridge University Press, 1987); M. Kaku, *Introduction to Superstrings* (Springer-Verlag, 1988).
2. D.J. Gross, J.A. Harvey, E. Martinec, and R. Rohm, *Nucl. Phys.* **B256**, 253 (1985).
3. P.C. Argyres and S.-H.H. Tye, *Fractional Superstrings with Space-Time Critical Dimensions Four and Six*, Cornell preprint CLNS 91/1068 (1991), to appear in *Phys. Rev. Lett.*
4. V.G. Knizhnik and A.B. Zamolodchikov, *Nucl. Phys.* **B247**, 83 (1984).
5. A.B. Zamolodchikov and V.A. Fateev, *Sov. Phys. J.E.T.P.* **62**, 215 (1985).
6. D. Gepner and Z. Qiu, *Nucl. Phys.* **B285**, 423 (1987).
7. D. Kastor, E. Martinec, and Z. Qiu, *Phys. Lett.* **200B**, 434 (1988); J. Bagger, D. Nemeschansky, and S. Yankielowicz, *Phys. Rev. Lett.* **60**, 389 (1988); F. Ravanini, *Mod. Phys. Lett.* **A3**, 397 (1988). See also S.-W. Chung, E. Lyman, and S.-H.H. Tye, *Fractional Supersymmetry and Minimal Coset Models in Conformal Field Theory*, Cornell preprint CLNS 91/1057 (1991).
8. P.C. Argyres, J. Grochocinski, and S.-H.H. Tye, *Nucl. Phys.* **B367**, 217 (1991).
9. V.G. Kač, *Adv. Math.* **35**, 264 (1980); *Bull. AMS* **3**, 1057 (1980); *ibid.*, *Adv. Math.*
10. H. Kawai, D.C. Lewellen, and S.-H.H. Tye, *Phys. Lett.* **185**, 197 (1987).
11. P.C. Argyres, K.R. Dienes, and S.-H.H. Tye, *Parafermion Characters*, Cornell preprint CLNS 91/37 = McGill/91-37 (to appear).
12. D. Sénéchal, *Phys. Rev.* **D39**, 3717 (1989); Cornell University (August 1990).
13. See, *e.g.*, K.R. Dienes, *Phys. Rev. Lett.* **68**, 57 (1992). More than 120 000 of such models have been constructed.
14. P.C. Argyres, E. Lyman, and S.-H.H. Tye, *Fractional Superstring*, Cornell preprint CLNS 91/1057 (to appear).
15. J. Distler and Z. Qiu, *Nucl. Phys.* **B338**, 509 (1990).